

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
4 March 2004 (04.03.2004)

PCT

(10) International Publication Number
WO 2004/018673 A2

(51) International Patent Classification⁷: C12N 15/10

LORENZ, Patrick [DE/DE]; Nibelungenstrasse 91,
64653 Lorsch (DE). ECK, Jürgen [DE/DE]; Siegfried-
strasse 138, 64646 Heppenheim (DE).

(21) International Application Number:
PCT/EP2003/009223

(74) Agent: VOSSIUS & PARTNER; Siebertstrasse 4, 81675
Munich (DE).

(22) International Filing Date: 20 August 2003 (20.08.2003)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(26) Publication Language: English

(30) Priority Data:
02018210.1 20 August 2002 (20.08.2002) EP

(71) Applicant (*for all designated States except US*):
B.R.A.I.N. BIOTECHNOLOGY RESEARCH AND
INFORMATION NETWORK AG [DE/DE]; Darm-
städter Strasse 34, 64673 Zwingenberg (DE).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): QUAISER, Achim
[DE/DE]; Schwarzwaldstrasse 31, 78549 Spaichingen
(DE). OCHSENREITER, Torsten [DE/US]; Josephine
Bay Paul Center for Comparative, Molecular Biology
and Evolution, The Marine Biological Laboratory, 7
MBL Street, Woods Hole, MA 02543 (US). TREUSCH,
Alexander, H. [DE/DE]; Ringstrasse 3, 64409 Messel
(DE). KLETZIN, Arnulf [DE/DE]; Kittlerstrasse 15,
64268 Darmstadt (DE). SCHLEPER, Christa [DE/DE];
Frankensteinerstrasse 155, 64297 Darmstadt (DE).

Published:

— without international search report and to be republished
upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: ISOLATION AND CLONING OF DNA FROM UNCULTIVATED ORGANISMS

(57) Abstract: The present invention relates to a device for the isolation and/or purification of nucleic acid molecules suitable to bind and/or inactivate inhibitors of the activity of reagents or enzymes used for DNA manipulation and to separate a plurality of nucleic acid molecules with respect to their size. Moreover, the invention relates to a method for the isolation of a nucleic acid molecule comprising applying a sample to the device of the invention wherein said nucleic acid molecule preferably represents a fraction of the metagenome of a given habitat. Furthermore, the invention relates to a method for the generation of at least one gene library comprising nucleic acid molecules isolated by the method of the invention and to a nucleic acid molecule isolated by the method of the invention.

WO 2004/018673 A2

ISOLATION AND CLONING OF DNA FROM UNCULTIVATED ORGANISMS

The present invention relates to a device for the isolation and/or purification of nucleic acid molecules suitable to bind and/or inactivate inhibitors of the activity of reagents or enzymes used for DNA manipulation and to separate a plurality of nucleic acid molecules with respect to their size. Moreover, the invention relates to a method for the isolation of a nucleic acid molecule comprising applying a sample to the device of the invention wherein said nucleic acid molecule preferably is part of a sample which represents a fraction of the metagenome of a given habitat. Furthermore, the invention relates to a method for the generation of at least one gene library comprising nucleic acid molecules isolated by the method of the invention and to a nucleic acid molecule isolated by the method of the invention and with the device of the present invention.

Several documents are cited throughout the text of this specification. The disclosure content of the documents cited herein (including any manufacture's specifications, instructions, etc.) is herewith incorporated by reference.

Enzymes are highly efficient biological catalysts. As such they are key players in environmentally friendly technical conversion processes of modern sustainable biotechnology.

Enzymes particularly from microbial sources are active ingredients in many processes of the textile, detergent, pulp- and paper, food and feed industries. In addition widespread stereoselective substrate recognition and conversion make enzymes particularly attractive for synthetic organic chemists in need of chiral specificity. A bottleneck in the development of innovative technical processes based on enzymes is the supply with suitable new biocatalysts. Owing to their phylogeny and physiological diversity microorganisms constitute the largest resource of natural genetic and enzymatic diversity. However the largest proportion of microorganisms

evades cultivation under laboratory conditions (Amann et. al. (1995). *Microbiol Rev* 59, 143-69). Classic microbiology relying on cultivation of pure strains to provide homogenous and defined systems for homologous enzyme production and to supply genomic DNA for recombinant expression strategies therefore inevitably fails to access the entire biosynthetic potential harboured in this enormous natural resource. The recent development of strategies to directly isolate and clone genomic DNA from non-cultivated microbial consortia opens up new dimensions of accessible enzymatic diversity (Rondon et. al. (2000). *Appl Environ Microbiol* 66, 2541-7). Fundamental work on the handling of DNA from non-cultivated microorganisms - the so-called metagenome - (Handelsman et. al. (1998) *Chem Biol* 5, R245-9) by Torsvik (Torsvik and Goksoyr (1978) *Soil Biology and Biochemistry* 10, 7-12), (Torsvik (1980) *Soil Biology and Biochemistry* 12, 15-21), Somerville (Somerville et. al. (1989) *Appl Environ Microbiol* 55(3), 548-554) and Schmidt (Schmidt et. al. (1991) *Journal of Bacteriology* 173, 4371-4378) showed that genomic DNA can be directly isolated from complex microbial assortments as present, inter alia, in plancton or soil. This DNA may be digested and cloned into suitable vectors for recombinant maintenance in heterologous hosts to generate screenable gene libraries. Such metagenome libraries were shown to be useful in the identification of novel genes from uncultivated organisms. The discovery of novel enzymes by screening of non-normalised metagenome libraries from planctonic and soil sources has been reported in the literature (Cottrell et. al. (1999) *Appl Environ Microbiol* 65, 2553-7), (Henne et. al. (1999) *Appl Environ Microbiol* 65, 3901-7; Henne et. al. (2000) *Appl Environ Microbiol* 66, 3113-3116); (US-patent No : 5,849,491); (Rondon et. al. (2000) *Appl Environ Microbiol* 66, 2541-7). The list of enzyme activities discovered in this way (lipase, esterase, amylase, nuclease, chitinase, xylanase) is still rather small. Importantly also more complex activities like the production of bioactive secondary metabolites requiring entire gene clusters for expression have been identified in metagenomic libraries (MacNeil et. al. (2001) *J Mol Microbiol Biotechnol* 3, 301-8) (Wang et. al. (2000) *Org Lett* 2, 2401-4) (Brady et. al. (2001) *Org Lett* 3, 1981-4). Secondary metabolites, like polyketides, are often produced by enzyme complexes encoded by assortments of genes covering in excess of 100 kbp of contiguous DNA (Schwecke et. al. (1995) *Proc Natl Acad Sci U S A* 92, 7839-43). The cloning of such

large fragments of environmental DNA is much more challenging than cloning smaller DNA fragments and is substantially facilitated by the current invention. Proprietary technology for the cloning particularly of normalised environmental DNA and the screening of libraries generated thereby is described in US patents US 6,280,926; US 6,054,267; US 6,057,103; US 6,001,574 and PCT applications WO99/45154; WO98/58085; WO99/10539.

DNA directly extracted from microbial consortia in the context of their natural substratum usually is contaminated with substances inhibiting standard enzymatic manipulations that are essentially required for cloning, analysis or amplification of nucleic acids carrying genetic information. In particular the efficiencies of DNA digestion with restriction enzymes (Tsai and Olson (1992) *Appl Environ Microbiol* 58, 2292-5), (Tebbe and Vahjen (1993) *Appl Environ Microbiol* 59, 2657-65), the polymerase chain reaction (PCR) (Zhou et. al. (1996) *Appl Environ Microbiol* 62, 316-22), DNA-DNA hybridisation and bacterial transformation with environmental DNA (Tebbe and Vahjen (1993) loc. cit.) are inversely correlated with natural substrate derived inhibitor concentrations. Besides inorganic inhibitors like heavy metal ions, there are polysaccharides and in particular humic and fulvic acids that act as the single most important sources of above mentioned inhibitions. Humic and fulvic acids are high molecular weight heterocyclic polyphenols mainly of plant origin with an affinity to polynucleotides and strongly protein denaturing properties ((Young et. al. (1993) *Appl Environ Microbiol* 59, 1972-1974); see appended figure 1).

Yet, the efficient removal of such inhibitors is a prerequisite for all enzymatic manipulations required, e.g., for cloning DNA, in particular environmental DNA, into suitable vectors. Several strategies have been pursued. Simple dilution of contaminated DNA to bring inhibitor concentrations below a critical threshold may be sufficient if the subsequent enzymatic manipulation is of suitable power to compensate for the concomitant reduction in target/substrate concentration. Surely such dilution will significantly curtail the efficiency of most subsequent molecular manipulations necessary for cloning following simple mass- action laws. The polymerase chain reaction (PCR), owing to its exponential amplification strategy is powerful enough to generate strong signals even from very low target numbers and

often reducing the amount of input environmental DNA (and inhibitors) in a reaction will substantially increase the amount of product achieved (Tsai and Olson (1992) loc. cit.). Gelfiltration of contaminated DNA raw extracts has been used to physically separate DNA from inhibitors based on size differences (Tsai and Olson (1992) loc. cit.), (Jackson et. al. (1997) *Applied and Environmental Microbiology* 63, 4993-4995), (Miller (2001) *J Microbiol Methods* 44, 49-58). Charge differences between DNA and inhibitors were exploited in strategies using ion-exchange chromatography purification (Tebbe and Vahjen, (1993) loc. cit.); (Straub et. al. (1995) *Water Science and Technology* 31, 311-315); (Smalla et. al. (1993) *J Appl Bacteriol*, 74, 78-85). In a different approach substances showing selective affinity towards polyphenols like soluble polyvinylpyrrolidone (PVP, figure 2, relative molecular weight 10'000-360'000 Da), insoluble polyvinylpolypyrrolidone (PVPP, a crosslinked derivative of PVP) or CTAB (hexadecyltrimethylammonium bromide) have been used to absorb (Holben et. al. (1988) *Appl. Environ. Microbiol.* 54, 703-711) or precipitate inhibitors from solutions (Zhou et. al. (1996) loc. cit.). Berthelet and co-workers used a PVPP affinity-matrix to chromatograph contaminated DNA solutions on spin columns (Berthelet et. al. (1996) *FEMS Microbiol Lett* 138, 17-22). Using ultracentrifugally generated CsCl density gradients Holben and co-workers (Holben et. al. (1988) loc. cit.) purified DNA from inhibitors based on equilibrium densities. For the construction of high quality libraries of uniform and particularly large DNA insert sizes (in vectors like Cosmid, Fosmid, BAC) a high resolution size selection step is essential to provide the reaction with uniformly sized insert DNA, especially if like in the case for BACs the cloning process does not feature any inherent size selective steps. This makes gel electrophoresis particularly attractive for the purification of environmental DNA. Hereby charge-mass ratios and size differences can be exploited simultaneously to achieve kinetic resolution of DNA from inhibitors and simultaneously the DNA itself is spread out according to size. Although simple gel electrophoresis may suffice to produce clonable DNA from soils containing only small amounts of humic and fulvic acids (Rondon et. al. (2000) loc. cit.), the humic content of soils varies greatly and can reach up to 60-80 % of the total organic matter (Tsai and Rochelle (2001) *Environmental Molecular Microbiology*, Horizon Scientific Press, page 15-30 (Extraction of nucleic acids from environmental samples)). Mostly therefore additional purification steps are necessary and still failures to produce

clonable soil DNA are common (Entcheva et. al. (2001) Appl Environ Microbiol 67, 89-99).

A particular modification of this method was devised by Young and co-workers (Young et. al. (1993) loc. cit.). Here, PVP was added to the gel to selectively lower the charge-mass ratio of humic acids so to improve resolution from DNA. This technique combines an affinity-based selective purification step retarding inhibitors in an electric field with a DNA size resolution step that is indispensable in the preparation of insert DNA for efficient large fragment cloning in vectors like BAC.

Yet, prior to subsequent enzymatic modifications of environmental DNA as required for cloning (like PCR using e.g. *Taq* or *Pfu* DNA polymerases or fill-up reactions using e.g. *Klenow*- or T4-DNA-polymerase or ligations using e.g. T4-DNA-ligase or multi-step reactions like phage-packaging) absorbants like PVP must be removed as they themselves are inhibitory for further enzymatic processes.

The separation of agarose gel-purified DNA from the PVP absorbant is achieved in the prior art by employing affinity chromatography after melting the DNA containing agarose slice (GeneClean® in (Young et. al. (1993) loc. cit.)). Such a procedure, however, is not suitable to purify very large DNA molecules because shearing forces generated during the elution process will cause fragmentation. Additionally elution efficiency is inversely correlated with molecule size, so large molecules will be selectively lost.

Alternatively, large DNA molecules can be electroeluted from an agarose slice cut from a gel after electrophoresis. The DNA will be recovered in diluted form in a buffer like TAE and has to be concentrated before further manipulation. This routinely involves precipitation in 70% ethanol (Ausubel et. al. (Eds.) (1998) Current Protocols in Molecular Biology, John Wiley & Sons, 2.11-2.1.10). Yet, such an alcohol precipitation involves at least one further centrifugation step and, accordingly, adverse shearing forces. Consequently, in order to purify large DNA fragments from agarose gels for enzymatic manipulation and cloning, prior art procedures involve melting a gel slice containing DNA, performing in-gel enzymatic manipulations in a re-solidified gel (like end polishing, ligation), solubilizing the gel using a degrading enzyme (Gelase® Epicentre, USA) and transforming the DNA into hosts (*E. coli*).

These procedures are complex and may lead to fragmentation or even loss of nucleic acid molecules. Furthermore, these manipulations of the prior art can not be carried out in the presence of enzyme inhibiting substances like PVP.

Many of the above strategies in different combinations have been used as part of multi-step purification protocols to produce clonable "metagenome DNA". Yet, environmental DNA purification is not trivial and whereas purification of DNA for PCR purposes may be accomplished using commercial kits (FastPrep® *Bio101*, USA), the preparation of sufficient amounts of concentrated and inhibitor-free high-molecular weight DNA (20-300 kbp) for cloning into Cosmids or BACs is much more challenging and may be doomed with failure (Entcheva et. al. (2001) loc. cit).

Thus, the technical problem underlying the present invention was to provide means and method for cloning of genetic material isolated from primary samples. The solution to this technical problem is achieved by providing the embodiments characterized in the claims.

The current invention provides means to overcome the technical difficulties associated with the isolation and cloning of large fragment DNA from uncultivated environmental sources.

Accordingly, the present invention relates to a device for the isolation and/or purification of nucleic acid molecules comprising at least two layers, a first layer being adapted to bind and/or inactivate inhibitors of the activity of reagents or enzymes used in nucleic acid manipulation and a second layer being adapted to separate a plurality of nucleic acid molecules with respect to their size.

The term "device" as employed herein is an arrangement/construction comprising, inter alia, gels, or gel chambers or columns as defined herein. Preferably, the gels, gel chambers or columns form the device of the present invention. The nucleic acid molecules to be isolated and/or purified are isolated and/or purified by passing them through the at least two layers of the device as defined herein.

The term "inhibitors of the activity of reagents or enzymes used in nucleic acid manipulation" describes substances comprised in samples of soil, aquatic samples

or samples of symbiotic/parasitic consortia which inhibit the activity of reagents or enzymes used in nucleic acid manipulation. Examples of said substances are described herein above and comprise inorganic inhibitors, like, e.g. heavy metal ions, organic inhibitors, like polysaccharides and in particular humic and fulvic acids. Humic and fulvic acids are high molecular weight heterocyclic polyphenols mainly of plant origin with an affinity to polynucleotides and strongly protein denaturing properties. The chemical structure of humic and fulvic acids is shown in appended figures 1A and 1B. Moreover, the chemical properties of said groups of molecules is described in detail by Stevenson (Humus chemistry: genesis, composition, reactions (1994) Wiley New York) and Buffle (Les substances humiques et leurs interactions avec les ions minéraux (1977) Conference Proceedings de la Commission d'Hydrologie Appliquée de A.G.H.T.M.. l'Université d'Orsay, 3-10).

The term "reagents used in nucleic acid manipulation" as used in this context comprises substances like metal ions (e.g. Mg^{2+} , Mn^{2+} , Ca^{2+}), (charged) inorganic and organic molecules required for enzymatic activity or for enzymatic co-factors, said co-factors themselves, or stabilizers. The term "enzymes used in nucleic acid manipulations" relates to enzymes like RNase(s), DNase(s), DNA-polymerase(s), ligase(s) or kinase(s) which are used for nucleic acid manipulation.

The term "nucleic acid manipulation" as used herein comprises standard methods known by the person skilled in the art. Said methods comprise DNA-engineering, such as cloning methods of nucleic acid molecules, the mutation of nucleotide sequences of nucleic acid molecules or amplification methods, like, e.g. PCR. Examples for said methods are described in the appended examples and in laboratory manuals, e.g. Sambrook et. al. (1989) *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York; Ausubel et. al. (1998), loc. cit.. In particular, the term "nucleic acid manipulation" relates to the manipulation of DNA or RNA and corresponding cloning techniques.

The term "layer" defines in accordance with the present invention a physical matrix which is characterized by its ability to separate samples containing nucleic acid

molecules and also characterized by its ability to separate, if desired, different nucleic acid molecules by their physiological properties, like size or overall charge. The recited first layer is adapted to characterized by its ability to bind and/or inactivate inhibitors described herein above. The ability of the first layer to bind and/or inactivate inhibitors may be achieved by the addition of compounds with sufficiently high binding affinity to the above described inhibitors so to retard their mobility in aqueous solutions and reduce their effective free concentration so to relieve nucleic acids migrating through the device of the invention from comigrating inhibitors.

The recited second layer is characterized by its ability to separate a plurality of nucleic acid molecules with respect to their size. Accordingly, said physical matrix may be a form of a physical matrix, suited to separate nucleic acid molecules based on molecular sieving, e.g. comprising gels or polymers.

According to a preferred embodiment of the device of invention said first layer is arranged above the second layer.

The term "above" defines the position of the first layer relative to the second layer and relative to the direction in which the samples migrate through the layers of the device. Accordingly, the nucleic acid molecule to be isolated and/or purified with the device of the invention is first contacted with the physical matrix of the first layer and afterwards contacted with the matrix of the second layer. Thus, the present invention comprises devices in which the first layer is horizontally above the second layer as well as devices in which the first layer is vertically above the second layer. Such a device is illustratively exemplified in the appended examples as a gel comprising two phases/layers. Accordingly, as illustrated in the appended figures and examples and described herein, this device may be arranged in form of a gel. Therefore, the device of the invention may be a device, wherein said first layer is a first phase of a gel and said second layer is a second phase of said gel. In its broadest sense, the term "phase" of a gel indicates that this phase has a different overall chemical constitution than a further (second) phase of said gel. For example, the difference in the overall chemical construction may be due to the presence of chemicals/compounds that bind or inactivate the above mentioned inhibitors of the activity of reagents or enzymes used in nucleic acid manipulations.

Preferably, the device of the present invention comprises a gel, wherein said gel is an agarose-gel or a polyacrylamid-gel.

Methods for the preparation of said gels are known by the person skilled in the art and are described in the appended examples and in standard laboratory manuals, e.g. Sambrook et. al. (1989), loc. cit., Cold Spring Harbor, New York; Ausubel et. al. (1998), loc. cit..

Preferably, the device of the invention comprises in said first layer polyvinylpyrrolidone (PVP), or polyvinylpolypyrrolidone (PVPP) or combinations thereof. As demonstrated in the appended examples PVP and PVPP are immobile components of said first layer or are characterized at least by a lower mobility compared to the nucleic acid molecules and to bind or interact with the above characterized "inhibitors of the activity of reagents or enzymes used in nucleic acid manipulation".

Further examples of corresponding components of said first layer (i.e. inactivators of the inhibitors defined herein above) are functional molecules like CTAB, EDTA, EGTA, cyclodextrins, proteins, (poly)peptides, nucleic acids immobilized or tethered on appropriate matrices acting as catcher molecules or ion-exchanger. These functional molecules may act by complexing ions (like EDTA, EGTA,), precipitating polysaccharides (like CTAB), binding small hydrophobic molecules or uncharged small molecules, like cyclodextrins, binding specific surface structures (like proteins, (poly)peptides and aptamers). Said proteins may comprise, e.g., antibodies and (poly)peptides directed against inhibitors. Furthermore, lectins are envisaged as inactivators of the inhibitors defined herein. An example for a nucleic acids acting as catcher molecules in accordance with the invention are RNA-aptamers.

The term "(poly)peptide" as used herein summarizes a group of molecules which comprise the group of peptides, consisting of up to 30 amino acids, as well as the group of polypeptides, consisting of more than 30 amino acids.

Above described low molecular weight compounds comprise electrically charged compounds (e.g. CTAB, EDTA) and uncharged small molecules (e.g. cyclodextrin). Said compounds are soluble in aqueous solutions and, thus, mobile due to rapid diffusion and in particular mobile in an electrical field. In order to achieve a lower

mobility of said compounds compared to the nucleic acid molecules, said compounds may be coupled to the physical matrices (e.g. chemically coupled). Compounds which are soluble in aqueous solutions but not electrically charged (e.g. PVP) and compounds which are non-soluble in aqueous solutions (e.g. PVPP) show a lower mobility compared to the nucleic acid molecules and thus do not essentially require to be coupled to the physical matrices.

More preferably, the device the invention is a device, wherein said second layer is substantially free of PVP, PVPP, CTAB, EDTA, EGTA, cyclodextrins, proteins, (poly)peptides, nucleic acids or ion-exchanger.

The term "substantially free" is understood in accordance with the invention to define a layer which does not contain the recited compounds in an effective amount which is detectable by standard methods. Said standard methods are known in the art and comprise MS (mass spectrometry), FT-IR (fourier transform infrared spectrometry), NMR (nuclear magnetic resonance) or HPLC (high performance liquid chromatography). Accordingly, the second layer does, most preferably, not contain any of the recited compounds as an essential/effective element.

According to a more preferred embodiment the device of the invention is electrically biased to enhance flow of (a) sample(s) through the layers.

As known by the person skilled in the art nucleic acid molecules are negatively charged due to the ribose-phosphate framework. Accordingly, nucleic acid molecules migrate in an electrical field from the cathode (-) to the anode (+).

Examples for devices which are electrically biased to enhance flow of (a) sample(s) are devices for gel electrophoresis. Devices with continuous as well as devices with discontinuous electrical fields are particularly comprised by the present invention. Accordingly, the electrical field can be discontinuous due to a gradient of a salt (buffer salt), a pulsed field comprising e.g. different angles. Again, an electrically biased device in accordance with the present invention is shown in the appended figures and illustrated in the appended examples.

The invention relates to a device, wherein said first layer preferably comprises sample loading means.

The term "sample loading means" defines in accordance with the invention means for placing the sample comprising nucleic acid molecules in the device of the invention. Examples for said means are sample slots in a gel, the surface of the matrix of a column or a valve for injecting a sample onto a column.

The isolation of nucleic acid molecules by using a device of the invention in the format of a horizontal agarose-gel is described in the appended example 1 and in the format of a single column is described in the appended example 4. Examples for the corresponding devices are shown in the appended figures 3 and 4.

In a more preferred embodiment of the invention said loading means are provided in an array in an upper portion of the first layer, defining an array of columns, each being capable of isolating nucleic acid molecules.

Hence, a device according to the preferred embodiment of the invention comprises more than one means for placing the sample. Thus, it is possible to isolate nucleic acid molecules from different samples in parallel. An example for loading means provided in an array is a gel comprising different loading slots and, therefore, different lanes (lines in which the samples are separated). An alternative example for said array are columns which are arranged in groups, e.g. in a frame. Said frame may comprise low numbers of columns (two to twelve) for the isolation of nucleic acid molecules from low numbers of samples. Also in accordance with the invention are frames comprising medium numbers of columns (twelve to 96) as well as frames with high numbers of columns (more than 96) which are suitable for high throughput screens (HTS).

According to an alternative embodiment of the invention said first layer of the device is arranged below the second layer.

The term "below" defines the position of the second layer relative to the first layer and relative to the direction in which the samples migrate through the layers of the device. The present invention comprises devices for the isolation and/or purification of nucleic acid molecules in which the nucleic acid molecules are first contacted with the physical matrix of the second layer to separate the molecules with respect to their size and subsequently contacted with the physical matrix of the first layer to bind and/or inactivate the above defined inhibitors.

Accordingly, the present invention comprises devices in which the second layer is horizontally above the first layer as well as devices in which the second layer is vertically above the first layer. Most preferred is in this context a device in form of a column, wherein said column comprises said first and said second layer.

An alternatively preferred embodiment of the invention relates to a device, wherein said second layer is a first phase of a column and said first layer is a second phase of said column. As pointed out herein above, the first layer comprises functional molecules capable of inactivating or binding inhibitors of the activity of reagents or enzymes used in nucleic acid manipulation.

It is also envisaged, in accordance with the invention, that an enhancement of the flow of (a) sample(s) through the layers of the device is, for example, achieved by gravity or by the pressure of the flow of a diluent. Examples for appropriate diluents are buffer solutions. The appended Examples show a device in form of a column as described herein.

Preferably, the device of the invention comprises in said first layer (in a column preferably the second phase) a matrix comprising PVP or PVPP or combinations thereof. As described herein above PVP and PVPP are characterized at least by a lower mobility compared to the nucleic acid molecules and to bind or interact with the above characterized "inhibitors of the activity of reagents or enzymes used in nucleic acid manipulation".

Further examples of corresponding components of said first layer (i.e. compounds capable of inactivating the inhibitors of nucleic acid manipulations) are EDTA, EGTA, CTAB, cyclodextrins, proteins, (poly)peptides, nucleic acids acting as catcher molecules or ion-exchangers. Said proteins may comprise, e.g., antibodies directed against inhibitors. Also lectins are envisaged as inactivators. An example for said nucleic acids acting as catcher molecules in accordance with the invention are RNA-aptamers.

More preferably the device the invention is a device, wherein said second layer (in a column preferably the first phase) is a matrix which is substantially free of PVP,

PVPP, CTAB, EDTA, EGTA, cyclodextrins, proteins, (poly)peptides, nucleic acids or ion-exchangers.

The term "substantially free" is defined herein above.

According to a further preferred embodiment of the invention, said matrix of said first and/or second layer is agarose, sepharose™, sephadex™, sephacryl™, BioGel™, superose™ or acrylamide.

Variations of the samples comprising the nucleic acid molecules to be isolated and/or purified with the device of the invention may require specific materials of the matrix of the first and/or the second layer. Said variations may concern the quality of the comprised nucleic acid molecules as well as the quality of the sample itself with respect to characteristic substances which may be contained. Accordingly, the matrix of the first and/or the second layer may be a specific matrix for gel-filtration which allows a molecular sieving of the nucleic acid molecules. Suitable media for matrices are characterized by a specific size of the pores which is known by the person skilled in the art and described in the instructions provided by the manufacturer of commercially available media. Said matrices comprise media for gel-filtration. Examples of said media comprise sepharose™ (e.g. sepharose2B, sepharose4B, sepharose6B), sephadex™ (e.g. sephadex G200, sephadex G150) sephacryl™ and superose™ as well as bio-gel™ P100 and bio-gel™ P200. Moreover, said media comprise agarose, polymers as e.g. dextrans and acylamid based-resins. In particular, media are preferred which are suitable for two-phases-columns. Further suitable materials are known to the person skilled in the art.

In a further preferred embodiment of the device of the invention said nucleic acid molecule is DNA or RNA. Most preferably, said DNA is genomic DNA (gDNA).

A particularly preferred embodiment of the invention is a device, wherein said nucleic acid molecule is derived from (micro)organisms of soil, sediments, water, for example sea water, or symbiotic/parasitic consortia.

The term "soil" defines in accordance with the invention the complex product of geological and biological processes acting on inorganic minerals and biomass deposited on the earth surface. It contains the majority of microbial biodiversity on

earth (Whitman et. al. (1998) Proc Natl Acad Sci USA, 95(12, 6578-83) acting to recycle and biomineralize organic matter and serves as a substratum to anchor and nourish higher plants.

In the appended examples the preparation of nucleic acid molecules isolated from soil is exemplarily described.

Examples for symbiotic/parasitic consortia in accordance with the present invention are consortia isolated e.g. from animal tissues or organs, e.g. from gut, stomach, intestines, like appendix or insect-, bird- and mammalian-intestinal tracts or -guts, comprising ruminant-gut. Also envisaged are animal tissues or organs from annelid(s), mollusc(s), sponge(s), cnidaria, arthropod(s), amphibian(s), fish or reptile(s). However, it is also envisaged that nucleic acids from parasitic consortia from human tissue, organs, sputum, faeces, sperm, blood, urine or other body fluids are isolated.

More preferably said (micro)organisms from which said nucleic acid molecules are derived from are (micro)organisms of aquatic plankton, animal tissues and organs as described herein above, microbial mats, clusters, sludge flocs, or biofilms.

(Micro)organisms of the "aquatic plankton" comprises bacterial plankton, archaeal plankton, viruses, phytoplankton as well as zooplankton. Said (micro)organisms are known as small organisms.

Biofilms are microbial assemblages on a surface in "aqueous environments" in which microbes are embedded in a hydrated polymeric matrix. This matrix acts like a kind of glue, holding the microbes together, attaching them to the surface and protecting them from detrimental external influences. They may contain several taxonomically distinct species (e.g. bacteria, fungi, algae, and protozoa) and may form on surfaces of diverse composition like e.g. metals, glass, plastics, tissue, mineral and soil particles.

Microbial mats and clusters are microbial assemblages/aggregates similar to biofilms in composition however not necessarily as firmly attached to solid surfaces.

According to a further preferred embodiment of the invention said (micro)organisms are isolated and/or purified as consortia of coexisting species. Preferably it is envisaged that said (micro)organisms are isolated and/or purified as consortia of

coexisting species without previous separation/purification of single microorganismic species.

Preferably said consortia of coexisting species comprise at least one organism that cannot proliferate indefinitely in an artificial setting (e.g. a synthetic or semisynthetic culture medium) in the absence of other species and/or in the absence of the substratum it is isolated from, and wherein said substratum contains the above defined inhibitors of the activity of reagents or enzymes used in nucleic acid manipulation.

To obtain the DNA of said microorganisms they may either be bulk-separated mechano-chemically from most of the surrounding matrix they are attached to or embedded or suspended in (like water, soil, sediments, organic debris of plant or animal origin) before lysis (indirect lysis) or they may be lysed and extracted directly i.e. still in the context of/attached to their surrounding physical matrix (soil, biofilm, floc, cluster) that contains a plurality of potential inhibitors of molecular, in particular enzymatic manipulation. Bulk-separation of microorganisms may be accomplished e.g. through mechanical agitation, ion-exchange resin mediated desorption (optionally facilitated through substances added to the extraction buffers used like detergents, salts) followed by an optional concentration step like filtration (for suspended plankton) or suspending and differential gravitational sedimentation in a liquid. In both instances total nucleic acids of mixed origin are isolated irrespective of taxonomic status, abundance and cultivability of the respective taxonomically mixed species.

Classic isolation of nucleic acid molecules from single microbial species or groups of microbial species comprise cultivation of said organisms by massive dilution in selectively growth supporting media. Thereby above mentioned inhibitors of molecular manipulation of nucleic acids are diluted to facilitate subsequent isolation and cloning. Yet at the cost of a massive reduction in species representation as very few species can be supported by standard cultivation techniques (see herein above). In contrast the present invention provides means for the isolation of nucleic acid molecules derived from taxonomically mixed (micro)organisms without cultivation in synthetic media. Since such steps of cultivation under laboratory conditions results in depletion or at least in significant dilution of organisms which do not grow under said

conditions the present invention surprisingly provides means for the isolation and/or purification of nucleic acid molecules derived from said organisms.

In a preferred embodiment of the invention said nucleic acid molecules represent a fraction of the metagenome of a given habitat.

As known in the art the term "metagenome" defines the totality of all genomes of organisms of a given habitat and is furthermore defined in the art; see inter alia Handelsman et. al. (1998) loc. cit.. In particular, the term "metagenome" relates to genomic nucleic acids, preferably DNA, derived from unknown or uncultivable microorganisms, i.e. organisms that cannot be isolated by standard methods and made actively replicating in standard artificial media for indefinite periods of time.

Accordingly the term "a fraction of the metagenome of a given habitat" defines in accordance with the invention nucleic acid molecules and in particular large nucleic acid molecules (>200bp) derived from the total pool of heterogenous microbial genomes present in a given habitat. This is irrespective of phylogenetic affiliation or molecular or physiological traits. Particularly the representation of any particular microbial genome in the extracted portion of the metagenome is not influenced by or dependent on the cultivatability of this organism. Therefore nucleic acids of uncultivated and in a preferred form particularly of uncultivable (micro)organisms are substantially represented in the extracted fraction of the metagenome.

An alternative embodiment of the invention relates to a method for the isolation of a nucleic acid molecule comprising applying a sample to the device as defined herein above. Said sample (for example derived from soil, sediments, water or symbiotic/parasitic consortia) is loaded onto a device comprising the above-identified at least two layers (for example a gel or a column comprising said two layers) and the nucleic acid molecule is purified and/or isolated by passing them through said two layers. In accordance with the invention, the first layer as defined herein, i.e. the layer comprising the inactivators of the inhibitors of reagents and enzymes of nucleic acid manipulations,4 retains said inhibitors, wherein said second layer provides for, e.g., an isolation and separation step for isolating/separating the nucleic acid molecules in accordance with their physical properties, like size or overall charge. It is envisaged and documented in the appended examples, that the nucleic acid

molecules to be isolated pass completely through the two layers of the device of the invention (e.g. a column comprising the two layers) or that the nucleic acid molecules pass only partially through the second layer. For example, it is envisaged that said nucleic acid molecules pass completely through said first layer comprising the inactivators of inhibitors and only partially through said second layer which is substantially free of said inactivators. This method is, inter alia, employed in the device of the invention which is in form of a gel. The nucleic acid molecules to be isolated and/or purified may be isolated or purified from the second layer by conventional means, for example by electroelution.

The method of the invention is also exemplified in the appended examples 1, 4 and 5.

Said method may optionally comprise one or more additional steps of subsequent purification of the obtained nucleic acid molecule(s).

According to a preferred embodiment of the method of the invention a fractions of the metagenome is isolated from a given habitat.

The invention relates in an alternative embodiment to a method for the generation of at least one gene library, comprising the steps of

- (a) isolating and/or purifying nucleic acid molecules from a sample using a device as defined herein above and optionally amplifying said nucleic acid molecules;
- (b) cloning the isolated and/or purified and optionally amplified nucleic acid molecules into appropriate vectors; and
- (c) transforming suitable hosts with said suitable vectors.

Methods for the amplification of an isolated and/or purified nucleic acid molecule are known in the art and comprise e.g. polymerase chain reaction (PCR).

Methods for the cloning of nucleic acid molecules into appropriate vectors and transforming suitable hosts with said suitable vectors represent standard methods of molecular biology and are described in the appended examples (in particular, example 2) and in laboratory manuals, e.g. Sambrook et. al. (1989) loc. cit.; Ausubel et. al. (1998), loc. cit.; Mülhardt (2002) Der Experimentator: Molekularbiologie/Genomics; Gustav Fischer. Suitable vectors are described herein below.

The above described method may, optionally, additionally comprise one or more of the following steps prior to the cloning of nucleic acid molecules into appropriate vectors according to step (b):

- (i) modification of DNA ends of the isolated and/or purified nucleic acid molecules, e.g. to remove or fill-up random 3'- or 5'-overhangs (polishing/fill-up/blunting), by DNA polymerase treatment (e.g. with Klenow enzyme, T4-polymerase) or treatment with exonucleases (like mung bean nuclease) or introducing defined 3'overhangs (like single nucleotide overhangs, in particular adenosin overhangs) using, e.g. DNA polymerase (like Taq), for subsequent cloning into appropriate vectors (e.g. T-overhang vectors like pGEM T-easy from Promega, USA) or introducing 3'- or 5'-overhangs using restriction endonucleases;
- (ii) phosphorylation of the isolated and/or purified nucleic acid molecules (e.g. by PNK); and/or
- (iii) ligation of the isolated and/or purified nucleic acid molecules to other nucleic acid molecules by treatment with an enzymatic ligase (e.g. by T4-ligase) or topoisomerase.

Also envisaged is a "sizing" step, as also illustrated briefly in (i) herein above, wherein said step comprises sizing the obtained nucleic acid molecules by treatment of the isolated and/or purified nucleic acid molecules with an enzymatic endonuclease (e.g. by restriction endonucleases or DNase I) and/or mechanical shearing (e.g. by ultrasonication or passing nucleic acids with high pressure through narrow tubes or valves similar to the "nebulizer" from Invitrogen, USA).

Suitable vectors comprise plasmid vectors (e.g. pUC18 and derivates thereof, pBluescript etc.), cosmid vectors (e.g. Expand, SuperCos), fosmid vectors (e.g. EpiFos 5), phage vectors (e.g. lambda ZAP), BAC vectors (e.g. pBeloBAC) and YAC vectors.

Preferably, said suitable hosts are selected from the group consisting of *E. coli*, *Pseudomonas* sp., *Bacillus* sp, *Streptomyces* sp, other actinomycetes, myxobacteria yeasts and filamentous fungi.

Transformation of said suitable host may be assayed by standard methods of molecular biology; see Sambrook et. al. (1989) loc. cit.; Ausubel et. al. (1998), loc.

cit.; Mülhardt (2002) loc. cit.. Corresponding assays for a successful transformation may be based on sequence similarity and performed on plated colonies (filter hybridization) by probe hybridization or by PCR analysis of colonies or extracted DNA of single or multiple colonies. Methods for the preparation of suitable oligonucleotides are known in the art. Alternatively the inserts may be sequenced and targets identified via homology search in appropriated sets of deposited sequence data (e.g. GenBank).

Activity based assays may be performed by screening for substrate conversion/degradation inside or outside the host (scoreable by substrate clearing zones around colonies; color development; molecular product profiling by high performance liquid chromatography (HPLC), mass spectrography (MS) or gas chromatography (GC); complementation of growth-deficient mutants), by screening for growth inhibition/stimulation of indicator organisms (in overlays).

Accordingly, and in a further embodiment, the present invention relates to a gene library obtained by the method disclosed herein and by employing the device described in this invention.

An alternative embodiment of the invention relates to a gene library generated from metagenome nucleic acid molecules, preferably from DNA, from non-planctonic (micro)organisms comprising average insert sizes of at least 50 kB, at least 55 kB, at least 60 kB, at least 70 kB, at least 80 kB, at least 90 kB or at least 100 kB.

As defined herein above planctonic (micro)organisms of the "aquatic plancton" comprises bacterial- and archaeal plancton, viruses, phytoplankton as well as zooplankton. Said (micro)organisms are known as small organisms living in aquatic habitats. Accordingly the term "non-planctonic (micro)organisms" defines (micro)organisms in accordance with the invention which are not comprised by the term "planctonic (micro)organisms". This group of (micro)organisms comprise (micro)organisms of soil, microbial mats, clusters sludge flocs, biofilms and symbiotic/parasitic consortia.

Similarly, the invention relates to a gene library generated from metagenome nucleic acid molecules, preferably DNA, from planctonic (micro)organisms comprising

average insert sizes of at least 85 kB, at least 90 kB, at least 95 kB, at least 100 kB, at least 120 kB, at least 140 kB, at least 160 kB or at least 200 kB.

In contrast to the group of "non-planctonic (micro)organisms" defined herein above the term "planctonic (micro)organisms" defines (micro)organisms of the "aquatic plancton" comprises bacterial- and archaeal plancton, viruses, phytoplankton and zooplankton as described herein above.

The average insert size of a (gene) library is, inter alia, determined by (a) isolating the cloned recombinant DNA of at least 0.1% of the clones of the respective library (however no less than a minimum of 20 clones) by methods known in the art, (b) digesting the isolated cloned recombinant DNA molecules with restriction enzymes (6-base or 8-base cutters, e.g., BamHI or EcoRI or Not I used singular or in combination) so to preferentially digest the vector backbone away from insert DNA (e.g. Not I used for pEpiFos5), (c) separating the resulting DNA fragments obtained from each clone individually by agarose gel electrophoresis (continuous or pulsed-field) as known in the art and (d) adding together the sizes of all non-vector fragments of all analyzed clones of a library and dividing the resulting number by the number of clones analyzed in order to obtain a figure for the average insert size.

The present invention relates further to a nucleic acid molecule comprising a DNA as depicted in SEQ ID NO: 1 or comprising a DNA as deposited under EMBL accession number AJ496176.

Said nucleic acid molecule of the invention has been isolated and obtained by employing the device of the invention. It represents a part of the genome of the newly identified Crenarchaeote as isolated with methods described herein and by techniques implied in the device of the present invention. Taxonomically the crenarchaeota represent a prokaryotic phylum as part of the archeal kingdom. The majority of its representatives are known to be hyperthermophiles yet increasingly they are found in mesophilic habitats as well; see Burggraf et al. (1997) Int J Syst Bacteriol., 47, 657-660; Preston et al. (1996) Proc Natl Acad Sci USA., 93, 6241-46. In the context of the present invention, the term "genome" defines not only sequences which are open reading frames (ORFs) encoding proteins, polypeptides

or peptides, but also refers to non-coding sequences. Accordingly, the term "nucleic acid molecule" comprises coding and, wherever applicable, non-coding sequences. The nucleic acid molecule of the invention furthermore comprises nucleic acid sequences which are degenerative to the above nucleic acid sequences. In accordance with the present invention, the term "nucleic acid molecule" comprises also any feasible derivative of a nucleic acid to which a nucleic acid probe may hybridize. Said nucleic acid probe itself may be a derivative of a nucleic acid molecule capable of hybridizing to said nucleic acid molecule or said derivative thereof. The term "nucleic acid molecule" further comprises peptide nucleic acids (PNAs) containing DNA analogs with amide backbone linkages (Nielsen, P., Science 254 (1991), 1497-1500). The term "ORF" ("open reading frame") which encodes a polypeptide, in connection with the present invention, is defined either by (a) the specific nucleotide sequences encoding the polypeptides specified above in (aa) or in (ab) or (b) by nucleic acid sequences hybridizing under stringent conditions to the complementary strand of the nucleotide sequences of (a) and encoding a polypeptide deviating from the polypeptide of (a) by one or more amino acid substitutions, deletions, duplications, insertions, recombinations, additions or inversions.

Furthermore the present invention relates in one embodiment to a nucleic acid molecule representing part of the genome of a non-thermophilic crenarchaeote, whereby said nucleic acid molecule has at least one of the following features:

- (a) it contains at least one ORF which encodes a polypeptide having the amino acid sequence SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21, SEQ ID NO: 23, SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 29, SEQ ID NO: 31, SEQ ID NO: 33, SEQ ID NO: 35;
- (b) comprises the DNA sequence of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, SEQ ID NO: 34;

- (c) it comprises portion of at least 20 nucleotides, preferably 100 nucleotides, more preferably at least 500 nucleotides which hybridize under stringent conditions to the complementary strand of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, SEQ ID NO: 34;
- (d) it is degenerate as a result of the genetic code with respect to the nucleic acid molecule of (c); or
- (e) it is at least 50% identical with the nucleic acid molecule of SEQ ID NO: 2, SEQ ID NO: 20 or SEQ ID NO: 30, 45% identical with the nucleic acid molecule of SEQ ID NO: 8 or SEQ ID NO: 26, 35% identical with the nucleic acid molecule of SEQ ID NO: 16, SEQ ID NO: 22 or SEQ ID NO: 24 or 30% identical with the nucleic acid molecule of SEQ ID NO: 4, SEQ ID NO: 14, SEQ ID NO: 18 or SEQ ID NO: 28;

The following list relates to the SEQ ID NOS as defined herein and shows (partial) identification of ORFs as defined herein:

Name	Identity
SEQ ID NO:1	complete DNA sequence 29i4, 33925 nt, of a newly identified Crenarchaeote
SEQ ID NO:2	ORF001 DNA sequence, 2367 nt, Fam. B DNA Polymerase (truncated ORF)
SEQ ID NO:3	ORF001 Protein sequence, 789 aa, Fam. B DNA Polymerase (truncated protein)
SEQ ID NO:4	ORF002 DNA sequence, 882 nt, α/β hydrolase
SEQ ID NO:5	ORF002 Protein sequence, 293 aa, α/β hydrolase
SEQ ID NO:6	ORF003 DNA sequence, 318 nt
SEQ ID NO:7	ORF003 Protein sequence, 105 aa
SEQ ID NO:8	ORF004 DNA sequence, 1086 nt, Polyhydroxyalkanoate Synthase
SEQ ID NO:9	ORF004 Protein sequence, 361 aa, Polyhydroxyalkanoate

	Synthase
SEQ ID NO:10	ORF005 DNA sequence, 582 nt
SEQ ID NO:11	ORF005 Protein sequence, 193 aa
SEQ ID NO:12	ORF006 DNA sequence, 438 nt
SEQ ID NO:13	ORF006 Protein sequence, 145 aa
SEQ ID NO:14	ORF007 DNA sequence, 915 nt, Glycosyl Transferase group 1
SEQ ID NO:15	ORF007 Protein sequence, 304 aa, Glycosyl Transferase group 1
SEQ ID NO:16	ORF008 DNA sequence, 1692 nt, Asparagine Synthase
SEQ ID NO:17	ORF008 Protein sequence, 563 aa, Asparagine Synthase
SEQ ID NO:18	ORF009 DNA sequence, 666 nt, Phosphoserin Phosphatase
SEQ ID NO:19	ORF009 Protein sequence, 221 aa, Phosphoserin Phosphatase
SEQ ID NO:20	ORF010 DNA sequence, 1212 nt
SEQ ID NO:21	ORF010 Protein sequence, 403 aa
SEQ ID NO:22	ORF011 DNA sequence, 1164 nt, Transmembrane protein
SEQ ID NO:23	ORF011 Protein sequence, 387 aa, Transmembrane protein
SEQ ID NO:24	ORF012 DNA sequence, 882 nt, Fix A Electron Transfer Flavoprotein
SEQ ID NO:25	ORF012 Protein sequence, 293 aa, Fix A Electron Transfer Flavoprotein
SEQ ID NO:26	ORF013 DNA sequence, 1284 nt, Fix B Electron Transfer Flavoprotein
SEQ ID NO:27	ORF013 Protein sequence, 427 aa, Fix B Electron Transfer Flavoprotein
SEQ ID NO:28	ORF014 DNA sequence, 1878 nt, Fix CX Fusion Electron Transfer Flavoprotein
SEQ ID NO:29	ORF014 Protein sequence, 625 aa, Fix CX Fusion Electron Transfer Flavoprotein
SEQ ID NO:30	ORF015 DNA sequence, 2238 nt, Sensory Transduction Histidin Kinase
SEQ ID NO:31	ORF015 Protein sequence, 745 aa, Sensory Transduction Histidin Kinase
SEQ ID NO:32	ORF016 DNA sequence, 519 nt

SEQ ID NO:33	ORF016 Protein sequence, 172 aa
SEQ ID NO:34	ORF017 DNA sequence, 1008 nt, (truncated ORF)
SEQ ID NO:35	ORF017 Protein sequence, 335 aa, (truncated protein)

A potential field of application of "ORF004" as defined herein comprise the generation or modification of biogenic polymers/polyesters (polyhydroxyalkanoate synthase, Zinn, (2001) Adv Drug Deliv Rev 53(1):5-21 and Fidler, (1992), FEMS Microbiol Rev 9(2-4):231-5; Snell, (2002) Metab. Eng. 4(1):29-40)

Furthermore, the ORF008, a potential asparagine synthetase may be used in the context of amino acid synthesis (EC 6.3.5.4) and/or for the generation of transgenic organisms, like bacteria, plants with altered capacities to generate amino acids.

In addition, ORFs 12, 13 or 14 may play a role in redox processes involved in nitrogen fixation and may be useful in generating transgenic organisms like bacteria, plants with altered capacities to assimilate nitrogen.

The term "hybridizing" as used herein refers to a pairing of polynucleotides to a complementary strand of polynucleotide which thereby form a hybrid. Said complementary strand polynucleotides are, e.g. the polynucleotides of the invention or parts thereof. Therefore, said polynucleotides may be useful as probes in Northern or Southern Blot analysis of RNA or DNA preparations, respectively, or can be used as oligonucleotide primers in PCR analysis dependent on their respective size. Preferably, said hybridizing polynucleotides comprise at least 10, more preferably at least 15 nucleotides in length while a hybridizing polynucleotide of the present invention to be used as a probe preferably comprises at least 100, more preferably at least 200, or most preferably at least 500 nucleotides in length.

It is well known in the art how to perform hybridization experiments with nucleic acid molecules, i.e. the person skilled in the art knows what hybridization conditions s/he has to use in accordance with the present invention. Such hybridization conditions are referred to in standard text books such as Sambrook et. al. (1989) loc. cit. or Higgins, S.J., Hames, D. "RNA Processing: A practical approach", Oxford University

Press (1994), Vol. 1 and 2.

"Stringent hybridization conditions" (also referred to highly stringent conditions as contrasted to conditions of low stringency) refers to conditions which comprise, e.g. an overnight incubation at 42°C in a solution comprising 50% formamide, 5x SSC (750 mM NaCl, 75 mM sodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1 x SSC at about 65°C. Said conditions for hybridization are also known by a person skilled in the art as "high stringent conditions for hybridization". Also contemplated are nucleic acid molecules that hybridize to the polynucleotides of the invention at lower stringency hybridization conditions ("low stringent conditions for hybridization"). Changes in the stringency of hybridization and signal detection are primarily accomplished through the manipulation of formamide concentration (lower percentages of formamide result in lowered stringency); salt conditions, or temperature. For example, lower stringency conditions include an overnight incubation at 37°C in a solution comprising 6X SSPE (20X SSPE = 3M NaCl; 0.2M NaH₂PO₄; 0.02M EDTA, pH 7.4), 0.5% SDS, 30% formamide, 100 µg/ml salmon sperm blocking DNA; followed by washes at 50°C with 1 X SSPE, 0.1% SDS. In addition, to achieve even lower stringency, washes performed following stringent hybridization can be done at higher salt concentrations (e.g. 5X SSC). Note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to suppress background in hybridization experiments. Typical blocking reagents include Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above, due to problems with compatibility.

Preferred in accordance with the present inventions are polynucleotides which are capable of hybridizing to the polynucleotides of the invention or parts thereof, under stringent hybridization conditions, i.e. which do not cross hybridize to unrelated polynucleotides.

The nucleic acid molecules that are homologous to the above-described molecules and that represent derivatives of these molecules usually are variations of these

molecules that represent modifications having the same biological function. They can be naturally occurring variations, for example sequences from other organisms, or mutations that can either occur naturally or that have been introduced by specific mutagenesis. Furthermore, the variations can be synthetically produced sequences. The allelic variants can be either naturally occurring variants or synthetically produced variants or variants produced by recombinant DNA processes.

Generally, by means of conventional molecular biological processes it is possible (see, e.g., Sambrook et. al. (1989) loc. cit.) to introduce different mutations into the nucleic acid molecules of the invention. One possibility is the production of deletion mutants in which nucleic acid molecules are produced by continuous deletions from the 5'- or 3'-terminus of the coding DNA sequence and that lead to the synthesis of proteins that are shortened accordingly. Another possibility is the introduction of single-point mutation at positions where a modification of the amino acid sequence influences, e.g., the enzyme activity or the regulation of the enzyme. By this method muteins can be produced, for example, that possess a modified K_m -value or that are no longer subject to the regulation mechanisms that normally exist in the cell, e.g. with regard to allosteric regulation or covalent modification. Such muteins may be identified, e.g. by methods of the present invention, to be valuable as therapeutically useful modulators (inhibitors/antagonists or enhancer/agonists) of the activity of the proteins of the present invention.

Nucleic acid molecules that hybridize to polynucleotides of the invention can be isolated, e.g., from genomic or cDNA libraries. In order to identify and isolate such nucleic acid molecules the polynucleotides of the invention or parts of these polynucleotides or the reverse complements of these polynucleotides can be used, for example by means of hybridization according to conventional methods (see, e.g., Sambrook (1989), loc. cit.). As a hybridization probe nucleic acid molecules can be used, for example, that have exactly or basically the nucleotide sequence of a part of the sequence shown in SEQ ID No: 1 or sequences complementary thereto. The fragments used as hybridization probe can be synthetic fragments that were produced by means of conventional synthesis methods and the sequence of which basically corresponds to the sequence of a nucleic acid molecule of the invention.

Preferably, the nucleic acid molecule of the invention is DNA or RNA.

An alternative embodiment of the invention relates to a vector comprising an above defined nucleotide acid molecule.

The vector of the present invention may be, e.g., a plasmid, phagemid, cosmid, fosmid, BAC, virus, bacteriophage or another vector used e.g. conventionally in genetic engineering, and may comprise further genes such as marker genes which allow for the selection of said vector in a suitable hosts and under suitable conditions.

Furthermore, the vector of the present invention may, in addition to the nucleic acid molecule of the invention, comprise expression control elements, allowing proper expression of the coding regions in suitable hosts. Such control elements are known to the artisan and may include a promoter, a splice cassette, translation initiation codon, translation and insertion site for introducing an insert into the vector. Preferably, the nucleic acid molecule of the invention is operably linked to said expression control sequences allowing expression in eukaryotic or prokaryotic cells.

Many suitable vectors are known to those skilled in molecular biology, the choice of which would depend on the function desired and include plasmids, phagemid, cosmids, fosmid, BAC, virus, bacteriophages and other vectors used conventionally in genetic engineering. Methods which are well known to those skilled in the art can be used to construct various plasmids and vectors; see, for example, the techniques described in Sambrook (1989) loc. cit. and Ausubel (1998) loc. cit.. Alternatively, the nucleic acid molecule and vectors of the invention can be reconstituted into liposomes for delivery to target cells. Thus, according to the invention relevant sequences can be transferred into expression vectors where expression of a particular (poly)peptide/protein is required. Typical cloning vectors include pBscpt sk, pGEM, pUC9, pBR322 and pGBT9. Typical expression vectors include pTRE, pCAL-n-EK, pESP-1, pOP13CAT. Typical prokaryotic cloning and expression vectors include: plasmid vectors like the pUC series (e.g. pUC18, pUC19), pGEM series (e.g. pGEM 7zf+, Promega, USA), pET series (e.g. pET22B, Novagen, USA), pBBC 1MCS series, pNOF (GL Biotech Germany), pCR-TOPO series and pCR Blunt (Invitrogen, USA), pBluescript series, pCAL series and pBC series (Stratagene, USA); Fosmid vectors like pEpifos5 and pCC1 (Epicentre, USA); Cosmid vectors like the Expand series (Expand I, II, III) (Roche, Germany), SuperCos (Stratagene, USA), pOJ436; BAC vectors like pBeloBAC, pCC1BAC (Epicentre, USA). However, the

present invention also envisages the expression of nucleic acid molecules as disclosed herein in eukaryotic vectors. Preferably, said nucleic acid molecules are linked to "control sequences". Said linking may be direct or indirect and refers, preferably to an operable linkage.

The term "control sequence" refers to regulatory DNA sequences which are necessary to effect the expression of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism. In prokaryotes, control sequences generally include promoter, ribosomal binding site, and terminators. In eukaryotes generally control sequences include promoters, terminators and, in some instances, enhancers, transactivators or transcription factors. The term "control sequence" is intended to include, at a minimum, all components the presence of which are necessary for expression, and may also include additional advantageous components.

The term "operably linked" refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences. In case the control sequence is a promoter, it is obvious for a skilled person that double-stranded nucleic acid is preferably used.

Thus, the vector of the invention is preferably an expression vector. An "expression vector" is a construct that can be used to transform a selected host cell and provides for expression of a coding sequence in the selected host. Expression vectors can for instance be cloning vectors, binary vectors or integrating vectors. Expression comprises transcription of the nucleic acid molecule preferably into a translatable mRNA. Regulatory elements ensuring expression in prokaryotes and/or eukaryotic cells are well known to those skilled in the art. In the case of eukaryotic cells they comprise normally promoters ensuring initiation of transcription and optionally poly-A signals ensuring termination of transcription and stabilization of the transcript. Possible regulatory elements permitting expression in prokaryotic hosts comprise, e.g., the PL, lac, trp or tac promoter in *E. coli*, and examples of regulatory elements permitting expression in eukaryotic host cells are the AOX1 or GAL1 promoter in yeast or the CMV-, SV40-, RSV-promoter (Rous sarcoma virus), CMV-enhancer, SV40-enhancer or a globin intron in mammalian and other animal cells. In this

context, suitable expression vectors are known in the art such as Okayama-Berg cDNA expression vector pcDV1 (Pharmacia), pCDM8, pRc/CMV, pcDNA1, pcDNA3 (In-vitrogen), pSPORT1 (GIBCO BRL). Typical prokaryotic cloning and expression vectors include: plasmid vectors like the pUC series (e.g. pUC18, pUC19), pGEM series (e.g. pGEM 7zf+, Promega, USA), pET series (e.g. pET22B, Novagen, USA), pBBC 1MCS series, pNOF (GL Biotech Germany), pCR-TOPO series and pCR Blunt (Invitrogen, USA), pBluescript series, pCAL series and pBC series (Stratagene, USA); Fosmid vectors like pEpifos5 and pCC1 (Epicentre, USA); Cosmid vectors like the Expand series (Expand I, II, III) (Roche, Germany), SuperCos (Stratagene, USA), pOJ436; BAC vectors like pBeloBAC, pCC1BAC (Epicentre, USA).

An alternative expression system which could be used to express a cell cycle interacting protein is an insect system. In one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. The coding sequence of a nucleic acid molecule of the invention may be cloned into a nonessential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of said coding sequence will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein coat. The recombinant viruses are then used to infect *S. frugiperda* cells or *Trichoplusia* larvae in which the protein of the invention is expressed (Smith, J. Virol. 46 (1983), 584; Engelhard, Proc. Nat. Acad. Sci. USA 91 (1994), 3224-3227).

In plants, promoters commonly used are the polyubiquitin promoter, and the actin promoter for ubiquitous expression. The termination signals usually employed are from the Nopaline Synthase promoter or from the CAMV 35S promoter. A plant translational enhancer often used is the TMV omega sequences, the inclusion of an intron (Intron-1 from the Shrunken gene of maize, for example) has been shown to increase expression levels by up to 100-fold. (Mait, Transgenic Research 6 (1997), 143-156; Ni, Plant Journal 7 (1995), 661-676). Additional regulatory elements may include transcriptional as well as translational enhancers. Advantageously, the above-described vectors of the invention comprises a selectable and/or scorable marker. Selectable marker genes useful for the selection of transformed cells and, e.g., plant tissue and plants are well known to those skilled in the art and comprise, for example, antimetabolite resistance as the basis of selection for dhfr, which

confers resistance to methotrexate (Reiss, Plant Physiol. (Life Sci. Adv.) 13 (1994), 143-149); npt, which confers resistance to the aminoglycosides neomycin, kanamycin and paromycin (Herrera-Estrella, EMBO J. 2 (1983), 987-995) and hygromycin (Marsh, Gene 32 (1984), 481-485). Additional selectable genes have been described, namely trpB, which allows cells to utilize indole in place of tryptophan; hisD, which allows cells to utilize histidinol in place of histidine (Hartman, Proc. Natl. Acad. Sci. USA 85 (1988), 8047); mannose-6-phosphate isomerase which allows cells to utilize mannose (WO 94/20627) and ODC (ornithine decarboxylase) which confers resistance to the ornithine decarboxylase inhibitor, 2-(difluoromethyl)-DL-ornithine, DFMO (McConlogue, 1987, In: Current Communications in Molecular Biology, Cold Spring Harbor Laboratory ed.) or deaminase from *Aspergillus terreus* which confers resistance to Blasticidin S (Tamura, Biosci. Biotechnol. Biochem. 59 (1995), 2336-2338).

Useful scorable markers are also known to those skilled in the art and are commercially available. Advantageously, said marker is a gene encoding luciferase (Giacomin, Pl. Sci. 116 (1996), 59-72; Scikantha, J. Bact. 178 (1996), 121), green fluorescent protein (Gerdes, FEBS Lett. 389 (1996), 44-47), β -glucuronidase (Jefferson, EMBO J. 6 (1987), 3901-3907) or secreted alkaline phosphatase (SEAP) (Schlatter et al. (2001) Biotechnol Bioeng. 5, 75(5), 597-606). This embodiment is particularly useful for simple and rapid screening of cells, tissues and organisms containing a vector of the invention.

The present invention furthermore relates to a host containing an aforementioned vector or an aforementioned nucleic acid molecule. Said host may be produced by introducing said vector or nucleotide sequence into the host by transfection or transformation wherein the nucleotide sequence and/or the encoded (poly)peptide/protein is foreign to the host. Upon the presence of said vector or nucleotide sequence in the host the expression of a protein encoded by the nucleotide sequence of the invention or comprising a nucleotide sequence or a vector according to the invention is mediated.

By "foreign" it is meant that the nucleotide sequence and/or the encoded (poly)peptide/protein is either heterologous with respect to the host, this means derived from a cell or organism with a different genomic background, or is

homologous with respect to the host but located in a different genomic environment than the naturally occurring counterpart of said nucleotide sequence. This means that, if the nucleotide sequence is homologous with respect to the host, it is not located in its natural location in the genome of said host, in particular it is surrounded by different genes. In this case the nucleotide sequence may be either under the control of its own promoter or under the control of a heterologous promoter. The vector or nucleotide sequence according to the invention which is present in the host may either be integrated into the genome of the host or it may be maintained in some form extrachromosomally. In this respect, it is also to be understood that the nucleotide sequence of the invention can be used to restore or create a mutant gene via homologous recombination.

Moreover, the present invention related to a method for producing a (poly)peptide as encoded by a nucleic acid molecule of the invention, comprising culturing the host of the invention under suitable conditions and isolating said (poly)peptide from the culture.

Isolation and purification of the recombinantly produced proteins and (poly)peptides may be carried out by conventional means including preparative chromatography and affinity and immunological separations involving affinity chromatography with monoclonal or polyclonal antibodies specifically interacting with said proteins/(poly)peptides. Preferably, said antibodies are antibodies of the invention as described herein below.

As used herein, the term „isolated protein“ includes proteins substantially free of other proteins, nucleic acids, lipids, carbohydrates or other materials with which it is naturally associated. Such proteins however not only comprise recombinantly produced proteins but include isolated naturally occurring proteins, synthetically produced proteins, or proteins produced by a combination of these methods. Means for preparing such proteins are well understood in the art. The proteins of the invention are preferably in a substantially purified form. A recombinantly produced version of said proteins, including secreted proteins, can be substantially purified by the one-step method described in Smith and Johnson, 1988.

An alternative embodiment of the invention relates to a (poly)peptide encoded by a nucleic acid molecule of the invention or as obtained by the method of the invention.

Preferably said (poly)peptide or fragment thereof is glycosylated, phosphorylated, amidated and/or myristylated.

Furthermore, the present invention relates to an antibody or an aptamer specifically recognizing the aforementioned (poly)peptide or a fragment or epitope thereof. Said antibody may be a monoclonal or a polyclonal antibody.

The term "fragment thereof" as used herein refers to fragments of said (poly)peptide/protein which are characterized by their capability to induce an immunological response in an immunized organism. Said response may be induced by the protein or fragment thereof either alone or in combination with a hapten, an adjuvant or other compounds known in the art to induce or elicit immunoresponses to a protein or fragment thereof.

The term "epitope" defines a single antigenic determinant. Said determinant is at least a portion of an antigen to which e.g. an antibody specifically binds to by its paratope; see Roitt et. al. (1993) Immunology 3rd edition, Mosby.

A preferred embodiment of the invention relates to an antibody which is a monoclonal antibody.

Said antibody, which is monoclonal antibody, polyclonal antibody, single chain antibody, or fragment thereof that specifically binds said peptide or polypeptide also including bispecific antibody, synthetic antibody, antibody fragment, such as Fab, a F(ab₂)', Fv or scFv fragments etc., or a chemically modified derivative of any of these (all comprised by the term "antibody"). Monoclonal antibodies can be prepared, for example, by the techniques as originally described in Köhler and Milstein, Nature 256 (1975), 495, and Galfré, Meth. Enzymol. 73 (1981), 3, which comprise the fusion of mouse myeloma cells to spleen cells derived from immunized mammals with modifications developed by the art. Furthermore, antibodies or fragments thereof to the aforementioned peptides can be obtained by using methods which are described, e.g., in Harlow and Lane "Antibodies, A Laboratory Manual", CSH Press, Cold Spring Harbor, 1988. When derivatives of said antibodies are obtained by the

phage display technique, surface plasmon resonance as employed in the BIAcore system can be used to increase the efficiency of phage antibodies which bind to an epitope of the peptide or polypeptide of the invention (Schier, Human Antibodies Hybridomas 7 (1996), 97-105; Malmberg, J. Immunol. Methods 183 (1995), 7-13). The production of chimeric antibodies is described, for example, in WO89/09622. A further source of antibodies to be utilized in accordance with the present invention are so-called xenogenic antibodies. The general principle for the production of xenogenic antibodies such as human antibodies in mice is described in, e.g., WO 91/10741, WO 94/02602, WO 96/34096 and WO 96/33735. Antibodies to be employed in accordance with the invention or their corresponding immunoglobulin chain(s) can be further modified using conventional techniques known in the art, for example, by using amino acid deletion(s), insertion(s), substitution(s), addition(s), and/or recombination(s) and/or any other modification(s) known in the art either alone or in combination. Methods for introducing such modifications in the DNA sequence underlying the amino acid sequence of an immunoglobulin chain are well known to the person skilled in the art; see, e.g., Sambrook (1989), loc. cit..

The term "monoclonal" or "polyclonal antibody" (see Harlow and Lane, (1988), loc. cit.) also relates to derivatives of said antibodies which retain or essentially retain their binding specificity. Whereas particularly preferred embodiments of said derivatives are specified further herein below, other preferred derivatives of such antibodies are chimeric antibodies comprising, for example, a mouse or rat variable region and a human constant region.

The term "scFv fragment" (single-chain Fv fragment) is well understood in the art and preferred due to its small size and the possibility to recombinantly produce such fragments.

The term "specifically binds" in connection with the antibody used in accordance with the present invention means that the antibody etc. does not or essentially does not cross-react with (poly)peptides of similar structures. Cross-reactivity of a panel of antibodies etc. under investigation may be tested, for example, by assessing binding of said panel of antibodies etc. under conventional conditions (see, e.g., Harlow and Lane, (1988), loc. cit.) to the (poly)peptide of interest as well as to a number of more or less (structurally and/or functionally) closely related (poly)peptides. Only those antibodies that bind to the (poly)peptide/protein of interest but do not or do not

essentially bind to any of the other (poly)peptides which are preferably expressed by the same organism/tissue as the (poly)peptide of interest, e.g. by a crenarchaeote, are considered specific for the (poly)peptide/protein of interest and selected for further studies in accordance with the method of the invention.

In a further alternative embodiment the present invention relates to a transgenic non-human mammal whose somatic and germ cells comprise at least one gene encoding a functional polypeptide selected from the group consisting of:

- (a) the polypeptide of the invention;
- (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and
- (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution.

A method for the production of a transgenic non-human animal, for example transgenic mouse, comprises introduction of the aforementioned polynucleotide or targeting vector into a germ cell, an embryonic cell, stem cell or an egg or a cell derived therefrom. The non-human animal can be used in accordance with the invention in a method for identification of compounds, described herein below. Production of transgenic embryos and screening of those can be performed, e.g., as described by A. L. Joyner Ed., *Gene Targeting, A Practical Approach* (1993), Oxford University Press. The DNA of the embryonal membranes of embryos can be analyzed using, e.g., Southern blots with an appropriate probe; see supra. A general method for making transgenic non-human animals is described in the art, see for example WO 94/24274. For making transgenic non-human organisms (which include homologously targeted non-human animals), embryonal stem cells (ES cells) are preferred. Murine ES cells, such as AB-1 line grown on mitotically inactive SNL76/7 cell feeder layers (McMahon and Bradley, *Cell* 62:1073-1085 (1990)) essentially as described (Robertson, E. J. (1987) in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*. E. J. Robertson, ed. (Oxford: IRL Press), p. 71-112) may be used for homologous gene targeting. Other suitable ES lines include, but are not limited to, the E14 line (Hooper et al., *Nature* 326:292-295 (1987)), the D3 line

(Doetschman et al., J. Embryol. Exp. Morph. 87:27-45 (1985)), the CCE line (Robertson et al., Nature 323:445-448 (1986)), the AK-7 line (Zhuang et al., Cell 77:875-884 (1994)). The success of generating a mouse line from ES cells bearing a specific targeted mutation depends on the pluripotency of the ES cells (i. e., their ability, once injected into a host developing embryo, such as a blastocyst or morula, to participate in embryogenesis and contribute to the germ cells of the resulting animal). The blastocysts containing the injected ES cells are allowed to develop in the uteri of pseudopregnant non-human females and are born, e.g. as chimeric mice. The resultant transgenic mice are chimeric for cells having either the recombinase or reporter loci and are backcrossed and screened for the presence of the correctly targeted transgene (s) by PCR or Southern blot analysis on tail biopsy DNA of offspring so as to identify transgenic mice heterozygous for either the recombinase or reporter locus/loci.

The transgenic non-human animals may, for example, be transgenic mice, rats, hamsters, dogs, monkeys, rabbits, pigs, or cows. Preferably, said transgenic non-human animal is a mouse.

The figures show:

Figure 1A shows the Structure of humic acids (after Stevenson 1982).

Figure 1B shows the Structure of fulvic acids (after Buffle 1977).

Figure 2 shows the Structure of polyvinylpyrrolidone (Monomer)

Figure 3 shows an example of an 2-phase PVP-low-melting agarosegel for simultaneous DNA purification and size resolution (Discontinuous Affinity-Gelelectrophoresis, DAG). The PVP-agarose-phase (the first phase) typically takes up approximately 1/4 to 1/3 of the gel but may take up maximally 80-95 % of the gel. After migrating through the PVP-phase into the agarose-only phase gel segments containing DNA fragments of interesting size may be excised and appropriately treated. The DNA migrates from minus (-) to plus (+).

Figure 4 shows an example of a 2-Phase Column chromatography for size resolution and affinity purification of DNA. In the first phase DNA in solution is resolved from inhibitors by size-exclusion chromatography. After passing this part of the composite column, the DNA passes a phase containing an affinity matrix (e.g. PVPP) to selectively bind and retard inhibitors. The DNA elutes from the column largest molecules first. The liquid flow is driven by hydrostatic or peristaltic pressure.

Figure 5 shows pulsed-field gelelectrophoretic (PFGE-) separations of metagenomic DNA isolated from soil (method A). In Figure 5 A the high molecular weight DNA is concentrated in a compression zone above 600 kbp. Yeast genomic DNA and a commercial size marker are added for reference. In Figure 5 B most DNA is sized to around 500 kbp but fragments up to 1.9 Mbp are visible.

M: Size marker (in kbp)

Figure 6 shows an insert analysis of metagenomic clones containing soil-derived DNA. Clones were digested with Not I.

Figure 7 shows an expression screening of arrayed fosmid clones containing metagenomic, soil-derived DNA. The encircled clone shows a halo of substrate degradation indicating hydrolase enzyme activity.

Figure 8 shows the quantification of metagenomic soil DNA in fractions eluting from columns after PVPP/sepharose 2B chromatography.

Figure 8 A: The DNA bands of fractions 6 - 22 were quantified by fluorescence intensity comparison after gel electrophoresis in a 1% agarose gel containing ethidium bromide. Gel documentation and analysis was done using GeneTools software from SynGene (UK). A 1% agarose gel was chosen to concentrate heterogeneous DNA fragments in a single band for the sake of simplifying quantification. This was achieved yet at the price of size-resolution. The apparent comigration of eluting DNA with the 23 kbp marker band therefore is an underestimation of true maximum DNA fragment sizes. Lane 1: λ Hind III Marker DNA

Figure 8 B: Analysis of DNA in PVPP/sepharose 2B chromatography fractions by agarose gel electrophoresis

* indicates fractions containing pure DNA

! indicates fractions containing large amounts of humic acids

Figure 9 shows the separation of soil metagenome DNA from humic substances by chromatography on PVPP/sepharose 2 B column.

Figure 9 A: Spectral absorption of eluting fractions 1–22 at 260 (DNA and humic acids) and 230 nm (humic acids) are plotted along their ratio and the relative DNA amounts as determined by agarose gel electrophoresis (see figure 8). A high A260/A230 ratio indicates pure DNA with low humic/fulvic acid contamination as can be seen in fractions 8 and 9. Absorption at 260 nm and 230 nm rises in two peaks in later fractions (12 and 17) possibly due to two size-populations of humic acids.

Figure 9 B: The result of a corresponding experiment is shown in Figure 9 B. Microbial DNA from humic substances in crude extract from soil is separated by PVPP/sepharose 2B chromatography. The gel electrophoretic analysis of fractions derived from this separation as described for Figure 9 A

Figure 10: Restriction digest (*NotI*) of randomly selected environmental fosmid clones separated by pulse field gel electrophoresis. Lanes 1,2: DNA size standards. A band of 7.5 kb visible in all lanes corresponds to the fosmid vector.

Figure 11: Phylogenetic tree based on full-length 23S rDNA sequences of Bacteria, Archaea and of sequences obtained from marine environmental genomic clones. Different alignment filters were used to evaluate the phylogenetic reconstruction. The tree topology shown here is based on a maximum parsimony analysis (using 1048 conserved positions selected by a positional variability filter). Closed circles indicate branching points supported by different phylogenetic methods and filters in reconstructions with the 23S and the corresponding 16S rDNA tree (see Experimental Procedures in appended example 5).

Figure 12: Schematic representation of the archaeal fosmid clone 29i4/SEQ ID NO: 1. Different shadings indicate the phylogenetic affinity of the putative protein-coding genes to archaea (diag. stripes), bacteria (dots), bacteria and archaea (vertical stripes), or archaea, bacteria and eukarya (grey). Hypothetical genes with no homologs are shown without fillings. ORF numbers match those in Table 1.

Figure 13: Phylogenetic analysis with selected sequences of the ETF-like protein family. Homologs of FixA proteins from archaea form a monophyletic group with the FixA proteins of nitrogen fixing bacteria and of *Thermotoga maritima*, clearly distinguished from "housekeeping" ETF proteins (Etf β homologs) of bacteria, archaea and eukaryotes. A third distinct subgroup (termed FixA paralog) is formed by a few as yet uncharacterized sequences from bacteria and archaea. For details of the phylogenetic reconstruction see Experimental Procedures in appended example 5.

Figure 14:

Figure 14 A: Size selection of soil DNA in a standard agarose gel

Preparative standard PFGE gel (EtBr stained) separation of soil DNA. The compression zone above 100 kbp was excised and resulting agarose plugs subsequently subjected to enzymatic digestion to test purity and clonability (Figure 14 B).

Figure 14 B: Failed hydrolyses of soil DNA (HindIII) purified in a standard agarose gel

Analytical PFGE showing DNA from an enzyme titration experiment (EtBr stained). DNA containing cubic plugs cut out from a standard PFGE agarose gel (Figure 14 A) and containing fragments sized over 100 kb were added to a reaction mix containing increasing units of *Hind*III restriction enzyme. After terminating the reaction the DNA was re-run on a second PFGE. It is evident from the similarly undigested DNA at 0U and at 100U enzyme concentration that this DNA is highly resistant to digestion with *Hind* III indicating inhibiting impurities that prevent molecular manipulation and cloning.

Figure 14 C: Size selection and purification of soil DNA in a 2-phase gel

Preparative 2-phase PFGE gel (SYBR green stained, inverted image) purification of soil DNA. The compression zone of DNA above 250 kbp was excised and subsequently subjected to enzymatic digestion as a proof of purity and clonability (Figure 14 D).

Figure 14 D: Successful partial hydrolyses of soil DNA (*Hind*III) purified in a 2-phase gel

Analytical PFGE showing soil DNA from an enzyme titration experiment (EtBr stained). 2-phase gel purified soil DNA cut out from a preparative 2-Phase Gel (Figure 14 C) and sized over 280 kb was *Hind* III digested and re-run on a second PFGE. It is evident that after purification in the 2-phase gel the DNA is readily digested to smaller sizes even with as little as 1U *Hind* III indicating a high degree of purity and access by enzymes necessary for subsequent molecular cloning.

Examples

The following examples illustrate the invention. These examples should not be construed as limiting: the examples are included for purposes of illustration and the present invention is limited only by the claims.

Example 1 Generation of a fosmid library from soil metagenomic DNA

1.1 Preparation of High Molecular Weight DNA from Soil (A)

Soil was collected from the upper layer of a partially ruderalized sandy ecosystem in Weiterstadt (Germany). About 50 g were suspended in 300 ml of buffer (pH 8, 20 mM Tris-HCl, 10 mM ϵ -aminocaproic acid, 10 mM EDTA) and incubated at 4°C for 15 h with gentle shaking. The suspension was sieved to remove larger particles and after centrifugation of the filtrate (5000 x g 30 min.) the resulting microbial fraction was embedded in agarose plugs (0.5 % low-melting SEAPlaque, FMC Bioproducts). The plugs were incubated at 37°C for 1 hour in lysozyme buffer (100 mM EDTA pH 8.0, 10 mM Tris-HCl pH 8.0, 50 mM NaCl, 0.2% Na-deoxycholate, 1% laurylsarcosin, 2 mg/ml lysozyme), then transferred into ESP solution containing 2 mg/ml proteinase K, 1% lauroyl Sarcosin and 0.5 M EDTA and incubated at 50°C for 24 h under gentle rotation and with 1 exchange of ESP buffer. 3 agarose plugs were placed in a 1%

agarose gel (Sigma A-2929) that contained 2 % PVP (Sigma PVP-360) in the upper part and no PVP in the lower part. Pulsed-field electrophoresis was performed in a CHEF-DR II PFGE machine (BioRad) at 10°C, 6 V cm⁻¹ for 20 h with 1 to 4 s pulses (Figure 5). A slice of agarose containing DNA in the size range of >30 kbp was cut out of the gel inserted into appropriately sized slots in a second gel and re-electrophoresed for a second size selection. After electroelution the resulting DNA was dialyzed and concentrated in a microconcentrator (Vivascience).

1.2 Preparation of High Molecular Weight DNA from Soil (B)

Alternatively total soil DNA was extracted using a protocol modified from Zhou and coworkers (Zhou et al., (1996) loc. cit.). Soil (5 g) was resuspended in 13.5 ml extractionbuffer (100 mM Tris/HCl pH8.0, 100 mM Na-EDTA, 100 mM Na-phosphate pH 8.0, 1.5 M NaCl, 1% CTAB) followed by an optional 3 cycles of freezing in liquid nitrogen and boiling in a microwave oven. After adding 1.5 ml of lysozyme solution (50 mg/ml) and 30 min. incubation at 37°C, 200µl proteinase K solution (10 mg/ml) were added followed by another 30 min. incubation at 37°C. Then 3 ml 10 % SDS was added followed by 2 hours incubation at 65°C. After centrifugation (10 min., 6000 x g, room temperature) the supernatant was collected and the pellet reextracted twice for 10 min. at 65°C with 4.5 ml of extraction buffer and 1 ml of 10% SDS. All supernatants were united and extracted with an equal volume of chloroform/isoamylalcohol (24:1 vol/vol). DNA was precipitated from the aqueous phase with 0.6 vol isopropanol, pelleted by centrifugation (16000 x g, 20 min., room temperature), washed in 70% ethanol and dissolved in 200µl TE buffer (10 mM Tris-HCl pH 8.0, 1 mM EDTA). Depending on the soil sample this solution was yellowish to dark brown. This DNA routinely defied any enzymatic manipulation by restriction enzymes and could be successfully used for PCR only at dilutions below 1/1000. It was therefore electrophoresed either in a constant voltage electrophoresis using a 0.5 % agarose 2-phase gel with 2 % PVP in the first phase (4 hours at 60 V) or in a 2-phase PFGE gel as described in (A).

Example 2: Library construction and analysis

About 0.5 µg of purified DNA was enzymatically treated to prepare 5' phosphorylated blunt ends and was ligated to the linearized and dephosphorylated fosmid vector pEpiFOS-5 (pEpiFOS Fosmid Library Production Kit, Epicentre). After in vitro packaging into lambda phages (Epicentre) and infection of E.coli strain EPI100 (Epicentre), cells were plated on LB medium containing 12.5 µg/ml chloramphenicol. The colonies were transferred to individual wells of 384-well microtitre plates containing 50 µl of LB with 12.5 µg/ml chloramphenicol and 7 % glycerol (v/v) and were incubated at 37°C for 24 hours. The library was stored at -80°C.

After blunt-end cloning into the fosmid vector pEpiFOS-5 about 50000 colonies were obtained per µg of soil DNA. Our final library was constructed from a single ligation mixture and contained 25278 clones arrayed in 66 384-well microtitre plates.

Restriction analysis of 30 randomly chosen clones with NotI showed insert sizes between 32.5 and 43.5 kbp, with an average of 36.5 kbp which corresponds well with the insert size range acceptable for this type of vector (Figure 6). The total library therefore contained an estimated 0.9 Gbp of environmental genomic soil DNA, which represents 225 genome equivalents assuming a 4 Mbp average genome size. Most inserts analyzed exhibited complex patterns after NotI digestion, suggesting that these clones contained DNA with high GC content.

Example 3: Screening of the library

The arrayed clones of the library were plated

- a) onto LB-agar to grow the cells for a subsequent preparation of fosmid-pools as a resource for sequence-based screenings (using PCR and degenerate primers to generate metagenome sequence tags e.g. as probes for hybridisation) and
- b) onto LB-agar containing specific substrates for the detection of enzymatically active colonies e.g. through scoring of clearing zones around the colonies (Figure 7) and
- c) onto LB-agar for growth and subsequent overlay with a lawn of indicator organisms to score for recombinants producing antibiotics.

Example 4: Purification of soil metagenomic DNA by use of a 2-phase gel-permeation/affinity column

Metagenome DNA from soil was extracted by gentle chemical lysis as described before (Zhou et al., (1996) *loc. cit.*). Crude DNA extract was passed over a 2-phase column by gravity-flow using a borosilicate glass column (BIORAD #737-0717). The column (7x150 mm) was packed with a lower phase of 1ml PVPP (SIGMA # P-6755) and an upper layer of Sepharose 2B (SIGMA # 2B-300) to a final volume of 5 ml. After equilibrating the column with 20ml of running buffer (100 mM NaCl, 10 mM Tris, 1 mM EDTA; pH 8.0) separation of DNA from humic and fulvic acids was initiated by applying crude DNA extract (ideally 1-5 % resin volume) to the top of the column. Eluting fractions of 300 µl were collected dropwise (6 drops) and subsequently analysed. Relative DNA content was quantified by comparing the fluorescence of DNA bands in each fraction after electrophoresis on a 1% agarose gel containing ethidium bromide (figure 8) and documentation using a CCD camera system (GeneGenius) and GeneTools software package (both Syngene, UK). The spectral absorption of fractions at 230 nm and 260 nm was measured spectrophotometrically. Fractions 8 and 9 contain pure clonable DNA. Analysis of later eluting fractions 10–12 show humic acid contamination as can be seen in the decrease in the ratio OD260/OD230. This is due to the significant rise of the OD230 values - the maximum absorption for humic acids. As humic substances also absorb at longer wavelengths the OD260 values increase similarly. The separation of fractions containing pure DNA and fractions containing humic substances can be improved further by adjusting the ratio of the loaded volume of crude DNA extract to the resin volume of the column in a way known to those skilled in the art and by adjusting the DNA content of the loaded samples.

Example 5: Exemplified identification of genes of a microorganism as part of a metagenome in accordance with the invention.

Traditionally, soil microbiology has focused on the description of cultivable microorganisms, while functional aspects of soil microbial communities have mainly been restricted to bulk studies that involved the monitoring of substrates and levels of end products. The application of molecular techniques to microbial ecology

revealed that many of the microbial transformations in the environments might be performed by organisms that have not yet been cultivated and thus far remained uncharacterized (Pace (1997) *Science* 276, 734-740, Hugenholtz et. al. (1998) *J. Bacteriol.* 180:4765-74). Soil was confirmed to be particularly rich in microbial diversity based on phylogenetic studies with 16S rRNA genes directly amplified from environmental samples (see e.g. Hugenholtz et. al. (1998) loc. cit., Borneman et al. (1996) *Appl. Environm. Microbiol.* 62, 1935-1943., Barns et. al. (1999) *Appl. Environm. Microbiol.* 65, 1731-1737, Dunbar et. al. (1999) *Appl. Environm. Microbiol.* 65, 1662-1669). Frequently, evidence was even found for the existence of microorganisms from divisions that were not previously associated with soil habitats and of which no cultivated relatives. Among one of the most striking discoveries was the frequent detection of non-thermophilic members of the archaeal kingdom Crenarchaeota (DeLong (1998) *Curr. Opin. Genet. Dev.* 8, 649-654). 16S rDNA sequences of these archaea were first identified in marine picoplankton (DeLong (1992) *Proc. Natl. Acad. Sci. U S A* 89, 5685-9, Fuhrman et. al. (1992) *Nature* 356, 148-9) and then found in freshwater habitats (Hershberger et. al. (1996) *Nature* 384, 420, Schleper et. al. (1997a) *Appl. Env. Microbiol.* 63, 321-323, McGregor et. al. (1997) *Appl. Env. Microbiol.* 63, 1178-1181, Jurgens et. al. (2000) *FEMS Microbiol. Ecol.* 34, 45-56) and in soils from various locations in the United States (Bintrim, et. al. (1997) *Proc. Natl. Acad. Sci. USA* 94, 277-282, Buckley et. al. (1998) *Appl. Environ. Microbiol.* 64, 4333-4339), Finland (Jurgens et. al. (1997) *Appl. Environ. Microbiol.* 63, 803-805), Japan (Ueda et. al. (1995) *Eur. J. Soil Sci.* 46, 415-421, Kudo et. al. (1997) *Biosci. Biotechnol. Biochem.* 61, 917-20) and Germany (Sandaa, et. al. (1999) *Appl. Environ. Microbiol.* 65, 3293-3297, Ochsenreiter et. al., in preparation). The ubiquitous ecological distribution of crenarchaeota was very surprising, because their cultivated relatives are exclusively (hyper)thermophiles isolated from terrestrial and marine hot springs.

Quantitative estimates have demonstrated the significant occurrence of non-thermophilic crenarchaeota in marine habitats (Massana et. al. (1997) *Appl. Environ. Microbiol.* 63, 50-6, Karner et. al. (2001) *Nature* 409, 507-510), in freshwater sediments (McGregor et. al. (1997) loc. cit.) and in soil (14,19). Some crenarchaeotal lineages were shown to be specifically associated with plant roots, indicating that the

organisms might play a role in the ecology of the rhizosphere (Simon et. al. (2000) *Environ. Microbiol.* 2, 495-505, Chelius & Triplett (2001) *Microb. Ecol.* 41,252-263). While 16S rRNA studies have provided insights into the huge extent of microbial diversity novel approaches are being sought to be able to functionally characterize those microorganisms that have escaped classical cultivation approaches.

Inspired by the rapid advances in microbial genomics of cultivated organisms, a novel approach has recently been initiated to characterize uncultivated organisms that have solely been predicted in rRNA gene surveys. It involves the construction of complex habitat-specific gene libraries by direct cloning of genomic fragments from environmental samples into cloning vectors (DeLong (2001) *Curr. Opin. Microbiol.* 4, 290-295). With the help of phylogenetically relevant gene markers, such as e.g. rDNA genes, large genomic fragments of specific phylotypes can be isolated from these libraries. Yet at present the full potential of this approach cannot be realized as technical constraints severely hamper the cloning of large DNA inserts, particularly from microbial consortia of inhibitor-rich environments like soils or sediments. The approach has successfully been applied to characterize uncultivated, marine microorganisms: Several genome fragments of the symbiotic crenarchaeote *Cenarchaeum symbiosum* and of marine archaea representing abundant components of the picoplankton in North Pacific and Antarctic waters were retrieved from BAC and fosmid libraries (Stein et. al. (1996) *J. Bacteriol.* 178,591-599, Schleper et. al. (1998) *J. Bacteriol.* 180, 5003-5009, Béjà et. al. (2000a) *Environ. Microbiol.* 2, 516-529, Beja et. al. (2002a) *Nature* 415:630-633, Béjà et. al. (2002b) *Appl. Environ. Microbiol.* 68, 335-45). A comparison of crenarchaeotal fosmids revealed significant genomic divergence even in clones with identical 16S rRNA sequences (Béjà et. al. (2002b) loc. cit.). The diversity of large photosynthetic gene clusters of proteobacteria was analyzed from marine planktonic genomic samples (Beja et. al. (2002a) loc. cit.). A novel type of rhodopsin, termed proteorhodopsin that functions as a light-driven proton pump was discovered in the genomic fragment of an uncultivated marine γ -proteobacterium (Béjà et. al. (2000b) *Science* 289, 1902-1906). The analyses of large genomic regions of hitherto uncultivated organisms also provided the basis for functional studies, including the monitoring of protein activities in the environment (Beja et. al. (2002a) loc. cit., Beja, et. al. (2001) *Nature* 411, 786-

9) and the characterization of proteins after expression in the surrogate host *E.coli* (Béjà et. al. (2000b) loc. cit., Schleper et. al. (1997b) *J. Bacteriol.* 179, 7803-7811).

The colocalization of functional, metabolic genes with phylogenetically ascribable genetic markers like rRNA genes provides insights into the physiological potential of uncultivated microorganisms. Clearly the likelihood of a physical colocalization of such markers on a contiguous cloned DNA stretch will be directly linked to DNA fragment size. This highlights the relevance of cloning large uninterrupted DNA fragments which is technically very difficult to achieve particularly from microbial consortia of inhibitor-rich environments like soils or sediments. We have developed procedures for the efficient purification of large DNA fragments by eliminating the polyphenolic compounds that heavily contaminate soil DNA. We have constructed complex genomic libraries and used these to isolate fragments from non-thermophilic crenarchaeota. While direct cloning of large DNA from soil samples has been demonstrated earlier (Rondon et. al. (2000) *Appl. Environ. Microbiol.* 66, 2541-2547), our study represents the first genomic characterization of a lineage of soil microorganisms that has solely been predicted by PCR-based studies.

5.1 Experimental Procedures

Preparation of DNA from Soil

Soil was collected from the upper layer (0 to 5 cm) of a partially ruderalized sandy ecosystem ("Am Rotböhl") near Darmstadt (Germany) in early Spring 2001. DNA was prepared as described in Example 1.1, supra. About 50 g were suspended in 300 ml of buffer (20 mM Tris pH 8, 10 mM ϵ -aminocaproic acid, 10 mM EDTA) and incubated at 4°C for 15h with gentle shaking. The sample was filtered to remove larger particles, and the microbial fraction was centrifuged and embedded into agarose plugs (0.5% low-melting SEAPlaque, FMC Bioproducts). These plugs were incubated at 37°C for 1 hour in 100 mM EDTA, 10 mM Tris-HCl pH 8, 50 mM NaCl, 0,2% deoxycholate, 1% lauroyl sarcosine, 1 mg/ml lysozyme, then transferred into ESP buffer (2 mg/ml proteinase K, 1% lauroyl sarcosine, 0.5 M EDTA) and incubated at 50°C for 24 h with gentle rotation and with one exchange of buffer. Agarose plugs were placed in a 1% agarose gel (Sigma A-2929) which contained 2% polyvinylpyrrolidone (VP-360, Sigma) in the first half and no PVP in the second half. Pulse field gel electrophoresis was performed at 10°C, 6 V cm⁻¹ for 20 h with 1 to 4

sec pulses in a CHEF-DR II (BioRad). DNA of > 30 kb was extracted from the gel and submitted to a second size selection using a regular agarose gel. After electroelution the resulting DNA was dialyzed and concentrated in a microconcentrator.

Library construction

Purified DNA (0.5 µg) was enzymatically treated to prepare 5' phosphorylated blunt ends and was ligated into fosmid vector pEpiFOS-5 (pEpiFOSTM, Epicentre). After *in vitro* packaging into lambda phages, the infected cells were plated on LB⁺ medium (containing 12.5 µg/ml chloramphenicol). The colonies were transferred to 384-well microtitre plates containing 50 µl of LB⁺ medium and 7 % glycerol (v/v). The plates were incubated at 37°C for 24 hours.

16S rDNA diversity studies

Primers specific for the domain Archaea (20F/958R, DeLong (1992) *Proc Natl Acad Sci USA* 89: 5685-9) and Bacteria (27F/1391R, Reysenbach and Pace (1995) *Archaea: a laboratory manual* (Cold spring Harbor Laboratory Press)) were used to amplify 16S rDNA fragments from the DNA used for constructing the large-insert library. The fragments were subsequently cloned into pGEM-T-easy (Promega) and sequenced. The ARB-software package (Ludwig et al. (1998) *Electrophoresis* 19: 554-568.) was used for alignments and phylogenetic analyses of the partial 16S rDNA genes.

Screening and sequence analysis of Fosmid clone 29i4

Plasmid DNA from the library was prepared from pools of 384 clones and screened by PCR with archaea-specific 16S rDNA primers (DeLong (1992) loc. cit.). A product of correct size (950 bp) was obtained from pool 29. It was randomly labelled with digoxigenin (Roche Biochemicals) and used as a probe in colony hybridization to identify the individual clone (i4). A subclone library was prepared from the fosmid DNA by mechanical shearing and cloning of 2-3 kbp fragments into pGEM-T-easy (Promega). The ends of the cloned DNA fragments were sequenced with vector primers using ABI3700 capillary sequencers. Remaining sequence gaps were closed by primer walking with sequence-derived oligonucleotides.

Sequence annotation

The ORF identification and automatic gene annotation were done with the help of the MAGPIE program package (Gaasterland and Sensen (1996) *Biochimie* 78: 302-310). The Wisconsin Package (Heidelberg Unix Sequence Analysis Server, HUSAR) was used for additional searches with GCGBLAST, identification of PFAM domains and transmembrane segments, for secondary structure prediction and peptide motifs. A tRNA gene was identified using the tRNA scan server (<http://www.genetics.wustl.edu/eddy/tRNAscan-SE/>). Multiple alignments were done with PILEUP and CLUSTAL and manually corrected in SEQLAB. The sequence was deposited in EMBL under the accession no. AJ496176.

Phylogenetic Analyses

The ARB-software package (Ludwig et al. (1998) loc. cit.) was used for alignments and phylogenetic analyses of full-length 16S and 23S rDNA genes from Archaea and from the marine environmental clones of a euryarchaeote 37F11 (Béjà et al. (2000a) loc. cit.), and of two crenarchaeota (4B7, Stein et al. (1996) loc. cit., Béjà et. al. (2002b) loc. cit.). The topologies of the 23S rDNA tree were evaluated using the maximum parsimony (parsimony interactive) and the distance matrix (Felsenstein correction) method with different alignment filters (gap-filter, positional variability filter, maximum frequency filter). The topologies of the corresponding 16S rDNA tree were evaluated using the same methods and in addition the maximum likelihood (fastDNAmI) method with different alignment filters as described above. Phylogenetic analysis of the putative FixA gene (ORF12/SEQ ID NO: 24) was performed using the protein parsimony (PROTPARS) and neighbor-joining (NEIGHBOR) programs from PHYLIP version 3.6 and PAUP version 3.1.1. The same overall topology was found with both methods based on 211 conserved positions from a sequence alignment of 33 FixA/ETFB (flavoprotein containing electron transport chain) homologs.

5.2 Results

Construction of a fosmid library from soil DNA

For preparation of high molecular weight DNA biomass from a soil sample was embedded into agarose plugs prior to lysis. The resulting DNA appeared heavily contaminated with polyphenolic compounds (i.e. humic and fulvic acids) as indicated by a dark brownish appearance. In the process of optimizing the subsequent

purification steps, a pulse field electrophoresis procedure was developed that involved an agarose gel with two phases. It allowed purification of the DNA from soil substances through a PVP (polyvinylpyrrolidone) containing phase. In a second phase the PVP was subsequently eliminated, while a first size selection of the DNA was achieved. Highly concentrated, pure and clonable DNA in the size-range of 30 to 100 kb was recovered in this one-step electrophoresis procedure, thereby minimizing shearing effects that tend to occur in repeated electrophoresis procedures. The approach has been successfully applied to different soil samples, like ruderal, agricultural and forest soils, for rapid preparation of pure and concentrated high molecular weight DNA (data not shown). After blunt-end cloning into the fosmid vector about 50,000 colonies were obtained per µg of soil DNA. Our final library contained 25,278 clones.

Restriction analysis of 30 randomly chosen clones using *NotI* showed insert sizes between 32.5 and 43.5 kb. The library therefore contained approximately 0.9 Gbp of environmental genomic soil DNA. 27 of 30 inserts analyzed exhibited complex patterns after *NotI* digestion, suggesting that these clones contained DNA with high G+C content (*NotI* recognition sequence: GCGGCCGC; Fig. 10).

Sequencing of insert ends from 2688 clones revealed in about 25 % of the sequences significant similarities to protein genes from the data bases (e-values of $< 10^{-10}$ in blastx searches). Among these were homologs of proteins from lineages that are typically found in soils, i.e. streptomycetes, clostridia and bacilli (data not shown). However most of the sequences did not show significant similarities to known protein genes. Together, these results confirmed that a great diversity of genomic DNAs was contained in the library.

To further monitor the diversity of the DNA used for construction of the library, a PCR-based 16S rDNA survey was performed using a primer set specific for Bacteria. 16S rRNA gene fragments affiliated with eight different bacterial phyla were identified in a random sample of 50 different sequences, many of which are typical for soil microbial assemblages, i.e. Actinobacteria, Chloroflexi, α, β, ϵ Proteobacteria, Planctomycetes, Acidobacterium/Holophaga, Cytophaga.

Identification and analysis of a genomic clone from non-thermophilic crenarchaeota in soil.

Using a multiplex PCR approach and 16S rDNA-specific probes, an archaeal fosmid clone was identified in the library. The insert of clone 29i4/SEQ ID NO: 1 was entirely sequenced and comprised 33,925 bp with an average G+C content of 40 %. It encoded a complete 16S and 23S ribosomal RNA operon, one tRNA gene and 17 predicted protein-encoding genes. The 16S RNA gene was 97 % identical over 711 positions (*E.coli* positions 8-719) to sequences previously recovered in a PCR study from the same soil (Ochsenreiter and Schleper, manuscript in preparation) and 95-97 % identical to sequences obtained from a soil in Wisconsin (Bintrim et al. (1997) loc. cit.). The ribosomal RNA operon of clone 29i4 consisted only of the 16S and 23S rRNA genes, without linked 5S rRNA or tRNA genes similar to cultivated thermophilic and uncultivated marine crenarchaeota. Phylogenetic analyses with the complete 16S rRNA and 23S rRNA genes confirmed the affiliation of clone 29i4 with the crenarchaeota (Fig. 11). As predicted by phylogenetic analyses with partial 16S rRNA sequences, it formed a sister group to the uncultivated marine organisms. The phylogenetic tree in Fig. 11 is based on complete 23S rRNA genes from known archaeal genomes and from environmental genomic fragments of marine archaea. The same branching orders were found in 16S rRNA phylogenies (see legend to Fig 11).

Ten of the 17 predicted protein-encoding genes showed significant similarity to genes of known function, two were conserved hypothetical genes, five open reading frames did not show any similarity to sequences in the databases (Febr. 2002). Eight of the predicted proteins showed highest similarities to archaeal homologs (Table 1 and Fig. 12). A family B DNA polymerase shared 46 % identical positions with its closest homolog from *Cenarchaeum symbiosum* (Schleper et al. (1997) J Bacteriol. 179, 7803-11). Although the C-terminal end of about 90 amino acids was not encoded on fosmid 29i4, all conserved exonuclease and polymerase motifs typically found in this class of DNA polymerases were identified in the deduced amino acid sequence.

Two other predicted proteins belonging to conserved archaeal protein families were an asparagine synthetase and a phosphoserine phosphatase, both involved in amino acid metabolism. Two putative glycosyl transferases (ORF07/SEQ ID NO.: 14 and 10/SEQ ID NO.: 20) shared significant similarities with homologs from the crenarchaeote *Sulfolobus solfataricus* and from the euryarchaeote *Pyrococcus* ssp.,

repectively. In contrast, a putative polyhydroxyalkanoate synthase (ORF04/SEQ ID NO.: 8) and a second α/β -hydrolase (ORF02/SEQ ID NO.: 4) were most closely related to bacterial proteins.

A gene cluster was identified with high similarity in structure and sequence to the fixABCX operons found in many symbiotic nitrogen-fixing soil bacteria. Based on its similarity to the components of the flavoprotein-containing electron transport chain (ETF) that is involved in β -oxidation of fatty acids in mitochondria and some bacteria, the operon was proposed to encode a flavoprotein-containing electron transport chain (Weidenhaupt et al. (1996) *Arch Microbiol* 165: 169-178). In symbiotic bacteria the operon is co-regulated with other genes involved in nitrogen fixation (Gubler and Hennecke (1998) *J Bacteriol* 170: 1205-1214). FixABCX genes have also been identified in the genomes of several other bacteria and some thermophilic and hyperthermophilic archaea, i.e. in *Sulfolobus solfataricus*, *Thermoplasma acidophilum*, *Pyrobaculum aerophilum* and *Aeropyrum pernix*. Phylogenetic analyses of the putative FixA gene from 29i4 and homologs from completely sequenced microbial genomes indicated a close affiliation of FixA from 29i4 with other archaeal proteins. Together with FixA proteins from nitrogen fixing bacteria they formed a distinct subgroup within the Etf β /FixA superfamily (Fig. 13).

A sensory histidine kinase was identified in close proximity but oriented in opposite direction to the fixABCX operon on clone 29i4. While the C-terminal half of the protein exhibited the conserved motifs typically found in sensory histidine kinases, no similarities to known proteins that might give hints to its specific role in sensing were found in the 350 amino-acid long N-terminal part.

Using PCR primers targeting the ends of the insert of clone 29i4 and internal protein coding genes, no contiguous genomic fragments overlapping clone 29i4 could thus far be detected in the library. However, additional archaeal clones were identified from non-thermophilic crenarchaeota in a second library that contained another 1.5 Gbp of DNA from the same soil sample. One of these clones was identified with archaea-specific 16S rRNA probes (as used for clone 29i4) and two other clones were identified through sequencing of insert ends from 768 randomly chosen clones. The sequence analysis of these genomic fragments is under way.

5.2 Discussion

The direct cloning of high-molecular weight DNA from soil is particularly difficult due to the occurrence of polyphenolic compounds that co-purify with DNA and severely inhibit PCR amplification reactions, hydrolysis by restriction enzymes, ligation and cloning procedures (Trevors and Van Elsas (1995) *Nucleic Acids in the Environment: Methods and Applications*, Springer Verlag, Berlin; Young et al. (1993) loc. cit.). Therefore, any purification protocol for DNA from soil samples must remove the phenolic compounds. Protocols have been developed that involve the addition of hexadecylmethylammoniumbromide (CTAB, Zhou et al. (1996) loc. cit.) or polyvinylpyrrolidone (PVP, Trevors and Van Elsas (1995) loc. cit.) in the extraction buffers, because these compounds complex polyphenolics or reduce their electrophoretic mobility when included in electrophoresis procedures (Young et al. (1993) loc. cit.). However, such compounds in turn have to be efficiently eliminated (e.g. by extraction, electrophoresis or affinity chromatography) because they inhibit enzymatic treatments of the isolated nucleic acids. Due to these difficulties most purification procedures either result in high quality DNA suitable for PCR amplification of gene fragments but too highly degraded for cloning of large fragments or it results in high molecular weight DNA that is not pure enough for cloning procedures. (refs above and own observations). Therefore, the device of the invention was developed for the electrophoresis procedure described herein above (two phases in which the DNA is first purified from polyphenolics in a PVP containing phase and subsequently cleaned in a second phase, thereby minimizing shearing effects that occur in repeated electrophoresis procedures). This technique allows reproducible obtainment of highly concentrated and pure, high molecular weight DNA. While purification techniques similar to those described by Zhou et al. (1996) loc. cit.) that was used by Rondon et al. (2000) loc. cit.) for cloning large DNA fragments were not applicable to many of our samples, the novel PVP electrophoresis procedure yielded reliable results with different soil samples of varying organic and humic acid content. Successful purification of the DNA was independent of the lysis procedures that we used, i.e. direct lysis of soil samples as in Zhou et al. (1996) loc. cit.) or lysis of microbial fractions as described here.

The complex environmental libraries constructed exemplarily by using the device and method of the invention contain a large fraction of the total genomic content of a soil microbial population, which has been referred to as the soil "metagenome" (Rondon et al. (2000) loc. cit.). The library characterized here was constructed in a BAC-derived fosmid containing cos-sites for *in vitro* packaging with lambda phages. Said vector yielded significantly larger clone numbers than using classical BAC-vectors (data not shown) and it allowed the direct cloning of undigested DNA by blunt-end ligation, thereby avoiding any bias introduced by restriction digests. Using 50 g of soil, a library with 0.9 Gbp of environmental DNA was constructed which represents approximately 225 genome equivalents assuming a 4 Mbp average genome size. Using archaea-specific 16S rDNA probes the fosmid clone 29i4 was identified. Sequence analysis demonstrated that a contiguous genomic fragment of non-thermophilic soil crenarchaeota was isolated: (i) Phylogenetic analyses based on the complete ribosomal 16S and 23S RNA genes indicated the specific affiliation with the crenarchaeotal clade as predicted in PCR-based studies. (ii) genes affiliated with archaea were found dispersed over the entire clone insert (iii) G+C content and codon usage of the predicted protein genes were similar over the entire insert and (iv) the deduced aminoacid sequence of a DNA polymerase gene showed greatest similarity to its homolog from the uncultivated marine symbiont *Cenarchaeum symbiosum*. Functional and biochemical analysis of the latter protein had confirmed the predicted non-thermophilic phenotype of this crenarchaeote (Schleper et al. 1997) J Bacteriol. 179, 7803-11).

On the other hand, significant differences to the content and structure of genomic fragments from uncultivated non-thermophilic marine archaea revealed that crenarchaeota from soil have significantly diverged from their relatives in other environments. An unusually large gap in the 16S-23S RNA operon and the lack of a GSAT gene (glutamate semialdehyde aminotransferase), which was consistently found to be directly linked to the ribosomal operon on all marine crenarchaeotal genome fragments (Béjà et al. (2002) loc. cit.) indicates the difference. Direct comparison of the soil fosmid clone to the genomic clones obtained from marine planctonic crenarchaeota and from the symbiont *C. symbiosum* revealed only one related protein encoding gene, i.e. the putative DNA polymerase. Genes on clone 29i4/SEQ ID NO: 1 appeared to be less densely packed than in genomes of other

archaea. Only 69 % of the sequence encoded RNA or protein genes. There was the large intergenic region in the 16S/23S rRNA cluster of 830 bp, which is atypical for ribosomal operons in crenarchaeota. Another large non-coding region of 2787 bp was found between ORF10/SEQ ID NO: 20 and fixA. No apparent genes or distinctive structural features, e.g. repetitive elements were identified in the non-coding regions.

The genomic information contained on fosmid 29i4/SEQ ID NO: 1 gives first insights into metabolic properties of crenarchaeota from soil and can serve as a basis for functional genomic studies. Beside genes encoding proteins for "house-keeping" functions (replication, aminoacid metabolism), two α/β type hydrolases so far seem to be particularly found in soil crenarchaeota. One of them encodes a putative protein involved in the synthesis of polyhydroxyalkanoates. The operon encoding FixABCX revealed a putative flavoprotein containing electron transport chain that is commonly found in symbiotic nitrogen-fixing bacteria. A detailed phylogenetic analysis indicated that the putative FixA protein of 29i4 is most closely affiliated with archaeal FixA homologs and not with the FixA proteins of Bacteria or the paralogous ETF proteins from other species of Archaea, Eucarya or Bacteria. Therefore, it seems unlikely that the fixABCX genes of 29i4 have been acquired by horizontal gene transfer e.g. from symbiotic nitrogen-fixing bacteria which might reside in the same soil habitat. They rather seem to originate from a common ancestor of the Archaea. None of the obligately aerobic archaea that contain the fixABCX genes is known to be capable of fixing nitrogen. Expression analysis of this operon in well-studied thermophilic model organisms, such as *Sulfolobus solfataricus* might shed light on its physiological role in crenarchaeota.

Example 6: Separation of DNA extracts on a PVPP/Sepharose 2B Column

Metagenome DNA from soil was extracted by gentle chemical lysis. Crude DNA extract was separated by gravity-flow gel filtration using a borosilicate glass column (BIORAD #737-0717). The column (7x150 mm) was packed with 1/6 volume PVPP (SIGMA # P-6755) and an upper layer of Sepharose 2B (SIGMA # 2B-300) to a final volume of 5 ml. Equilibration and removal of storage buffer was done by passing 5 reservoir volumes of TEN buffer (100 mM NaCl, 10 mM Tris, 1 mM EDTA; pH 8.0) by

gravity flow. Separation of DNA and humic acids was initiated by applying crude DNA extract (ideally 1-5 % resin volume) to the column. Approximately 300 µl fractions were collected dropwise (6 drops per fraction) and subsequently analysed. Relative DNA contents was quantified by fluorescence intensity comparison of DNA bands of each fraction on 1% agarose. Absorption at 230 nm and 260 nm was measured spectrophotometrically. Fractions 7 -10 represent pure DNA. Analysis of later fractions 11 – 22 show humic acid content with a maximum in fraction 16. As humic substances also absorb at longer wavelengths the OD₂₆₀ values increase in a similar extent as those of OD₂₃₀ which could mislead to assume high DNA concentrations. The specialist is able to discriminate pure DNA fractions from those contaminated by humic acids by increasing absorption at 230 nm and a low ratio OD₂₆₀/OD₂₃₀.

Example 7: Validation of the purifying effect of the 2-phase gel on high molecular weight soil DNA

Pellets of uncultivated bacteria isolated from soil were embedded in agarose plugs and treated appropriately to liberate their genomic DNA in situ. The DNA plugs were embedded either in an ordinary preparative PFGE (pulsed field gel electrophoresis) gel (Fig. 14 A) or a 2-phase PFGE gel (Fig. 14 C) for sizing and eventual purification. After completion of the PFGE, agarose plugs containing high molecular weight DNA in the compression zones were cut out and subjected to digestion with the restriction endonuclease *Hind* III to create DNA ends compatible with common cloning sites in suitable DNA vectors. After terminating the digestion the DNA plugs were inserted into analytical PFGE gels to analyse the success of the digestion.

As seen in Fig. 14 B soil DNA separated in a conventional agarose PFGE gel was highly resistant to digestion even with large amounts of restriction enzyme whereas high molecular weight DNA purified in the 2-phase PFGE gel was readily fragmented with very little restriction enzyme (Fig. 14D) proving its high degree of purity and consequently clonability. Cloning DNA fragments with compatible ends into suitably prepared DNA vectors like BAC, Plasmid, Lambda, Fosmid, Cosmid, YAC and PAC are standard procedures known to those skilled in the art.

Table 1:

Predicted RNA and protein encoding genes in the archaeal fosmid 29i4.

ORF	Nt range	protein size	putative function	most similar ortholog*	Phyl. aff.#	Comments
01	1-2367	789 aa (C-term. truncated)	Family B DNA polymerase pfam00135	AAC62689 Cenarchaeum symbiosum (0.0)	AEB	
RNA	3548-5018		16S RNA	85% identity to 16S RNA of Cenarchaeum symbiosum		
RNA	5847-8818		23S RNA	77% identity to 23S RNA of Cenarchaeum symbiosum		
02	8968-9849	293 aa	α/β hydrolase pfam00561	AAD02150 Pseudomonas stutzeri (e -17)	AEB	Catalytic triade Ser/Asp/His; close homologs are from bacteria
03	9888-10205	105 aa	Hypothetical	none		
04	10219-11304	361 aa	PHA Synthase (α/β hydrolase, pfam00561)	P45366 Thiocystis violacea (e -66)	B	No homolog in archaea but PHB production has been described in Halobacteriaceae
05	11285-11866	193 aa	Hypothetical	none		
RNA	12414-12502		tRNA ^{Arg}			
06	13017-13454	145 aa	Hypothetical	none		
07	14324-15238	304 aa	Glycosyl transferases group 1, pfam00534	AAK41834 Sulfolobus solfataricus (e -11)	AEB	Transfer of ADP, UDP, GDP, CMP linked sugars
08	15716-17407	563 aa	Asparagine synthetase pfam 00310	AAB99117 Methanococcus janaschii (e -67)	AEB	
09	17492-18157	221 aa	Phosphoserin phosphatase pfam00702	AAB86099 Methanothermobacter thermautotrophicus (e -14)	AEB	
10	18377-19588	403 aa	Conserved hypothetical	CAB50138 Pyrococcus abyssi (e -91)	A	Homologs only found in P. abyssi, P. horikoshii, A. pernix, Glycosyltransferase group 2 domain; pfam 00535
11	20630-21793	387 aa	Transmembrane protein	BAB50489 Mesorhizobium loti (e -31)	B	Domain pfam 01173
12	24580-25461	293 aa	Fix A pfam01012	P53576 Azotobacter vinelandii (e -23)	AB	Paralogs of ETF β : electron transfer flavoprotein β subunit
13	25458-26741	427 aa	Fix B pfam00766	P53578 Clostridium saccharobutylicum (e -45)	AB	Paralogs of ETF α : electron transfer flavoprotein α subunit
14	26738-28615	625 aa	Fix CX	NP_454687 Salmonella enterica (e -36) FixX: Thermoplasma volcanium (e -08)	AB	Fused protein of FixC and FixX
15	29228-31465	745 aa	Sensory transduction histidine kinase, pfam00512	BAB73503 Nostoc sp. PCC 7120 (e -12)	AEB	
16	32505-33023	172 aa	Hypothetical protein	none		
17	32918-33925	335 aa (truncated)	hypothetical	none		

*proteins are designated by their gene identification numbers followed by the species name. The e-values of blastx searches are added in brackets.

Phyl. aff. = Phylogenetic affinity, denotes occurrence of homologs in Archaea (A), Bacteria (B) or Eucarya (E).

Claims

1. A device for the isolation and/or purification of nucleic acid molecules comprising at least two layers, a first layer being adapted to bind and/or inactivate inhibitors of the activity of reagents or enzymes used in nucleic acid manipulation and a second layer being adapted to separate a plurality of nucleic acid molecules with respect to their size.
2. The device of claim 1, wherein said first layer is arranged above the second layer.
3. The device of claim 1 or 2, wherein said first layer is a first phase of a gel and said second layer is a second phase of said gel.
4. The device of claim 3, wherein said gel is an agarose-gel or a polyacrylamid-gel.
5. The device of any one of claims 1 to 4, wherein said first layer comprises polyvinylpyrrolidone (PVP), polyvinylpolypyrrolidone (PVPP), CTAB, EDTA, EGTA, cyclodextrins, proteins, (poly)peptides, antibodies, aptamers, lectins, nucleic acids or ion-exchanger.
6. The device of any one of claims 1 to 5, wherein said second layer is substantially free of PVP, PVPP, CTAB, EDTA, EGTA, cyclodextrins, proteins, (poly)peptides, aptamers, antibodies, lectins, nucleic acids or ion-exchanger.
7. The device of any one of claims 3 to 6, wherein the device is electrically biased to enhance flow of (a) sample(s) through the layers.
8. The device of any one of claims 3 to 7, wherein said first layer comprises sample loading means.

9. The device of claim 8, wherein said loading means are provided in an array in an upper portion of the first layer, defining an array of columns, each being capable of isolating nucleic acid molecules.
10. The device of any one of claims 1, 5, 6, 7, 8 or 9, wherein said first layer is arranged below the second layer.
11. The device of claim 10, which is a column comprising said first and said second layer.
12. The device of claim 10 or 11, wherein said second layer is a first phase of a column and said first layer is a second phase of said column.
13. The device of any one of claims 10 to 12, said first layer is a matrix comprising PVP, PVPP, CTAB, EDTA, EGTA, cyclodextrins, proteins, (poly)peptides, aptamers, antibodies, lectins, nucleic acids or ion-exchanger.
14. The device of any one of claims 10 to 13, wherein said second layer is a matrix which is substantially free of PVPP PVP, PVPP, CTAB, EDTA, EGTA, cyclodextrins, proteins, (poly)peptides, aptamers, antibodies, lectins, nucleic acids or ion-exchanger.
15. The device of claims 13 or 14, wherein said matrix of said first and/or second layer is selected from the group consisting of agarose, sepharoseTM, sephadexTM, sephacrylTM, BioGelTM, superoseTM and acrylamid.
16. The device of any one of claims 1 to 15, wherein said nucleic acid molecule is DNA or RNA.
17. The device of claim 16, wherein said DNA is genomic DNA.
18. The device of claim 16 or 17, wherein said nucleic acid molecule is derived from (micro)organisms of soil, sediments, water or symbiotic/parasitic

consortia.

19. The device of claim 18, wherein said (micro)organisms are (micro)organisms of aquatic plancton, microbial mats, clusters, sludge flocs, or biofilms.
20. The device of claim 18 or 19, wherein said (micro)organism are isolated as consortia of coexisting species.
21. The device of any one of claims 1 to 20, wherein said nucleic acid molecules represent a fraction of the metagenome of a given habitat.
22. A method for the isolation of a nucleic acid molecule comprising applying a sample to the device as defined in any one of claims 1 to 21.
23. The method of claim 22, wherein a fraction of the metagenome is isolated from a given habitat.
24. A method for the generation of at least one gene library, comprising the steps of
 - (a) isolating and/or purifying nucleic acid molecules from a sample using a device as defined in any one of claims 1 to 21 and optionally amplifying said nucleic acid molecules;
 - (b) cloning the isolated and/or purified and optionally amplified nucleic acid molecules into appropriate vectors; and
 - (c) transforming suitable hosts with said suitable vectors.
25. The method of claim 24, wherein said suitable hosts are selected from the group consisting of *E. coli*, *Pseudomonas* sp., *Bacillus* sp, *Streptomyces* sp, other actinomycetes, myxobacteria, yeasts and filamentous fungi.
26. A gene library obtained by the method of claim 24 or 25.
27. A gene library generated from metagenomic nucleic acid molecules from non-

planctonic (micro)organisms comprising average insert sizes of at least 50 kB, of at least 55 kB, of at least 60 kB, of at least 80 kB, of at least 90 kB or of at least 100 kB.

28. A gene library generated from metagenomic nucleic acid molecules from planctonic (micro)organisms comprising average insert sizes of at least 85 kB, at least 90 kB, at least 95 kB, at least 100 kB, at least 120 kB, at least 140 kB, at least 160 kB or at least 200 kB.
29. A nucleic acid molecule comprising a DNA as depicted in SEQ ID NO: 1 or a DNA as deposited under EMBL accession number A3496176.
30. A nucleic acid molecule representing a part of the genome of a non-thermophilic crenarchaeota, whereby said nucleic acid molecule has at least one of the following features:
 - (a) it contains at least one ORF which encodes a polypeptide having the amino acid sequence SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21, SEQ ID NO: 23, SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 29, SEQ ID NO: 31, SEQ ID NO: 33, SEQ ID NO: 35;
 - (b) comprises the DNA sequence of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, SEQ ID NO: 34;
 - (c) it comprises portion of at least 20 nucleotides, preferably 100 nucleotides, more preferably at least 500 nucleotides which hybridise under stringent conditions to the complementary strand of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, SEQ ID NO: 34;

- (d) it is degenerate as a result of the genetic code with respect to the nucleic acid molecule of (c); or
 - (e) it is at least 50% identical with the nucleic acid molecule of SEQ ID NO: 2, SEQ ID NO: 20 or SEQ ID NO: 30, 45% identical with the nucleic acid molecule of SEQ ID NO: 8 or SEQ ID NO: 26, 35% identical with the nucleic acid molecule of SEQ ID NO: 16, SEQ ID NO: 22 or SEQ ID NO: 24 or 30% identical with the nucleic acid molecule of SEQ ID NO: 4, SEQ ID NO: 14, SEQ ID NO: 18 or SEQ ID NO: 28;
- 31. The nucleic acid molecule of claim 29 or 30, which is DNA or RNA.
 - 32. A vector comprising the nucleotide acid molecule of any one of claims 29 to 31.
 - 33. A host transfected or transformed with the vector of claim 32.
 - 34. A method for producing a (poly)peptide as encoded by a nucleic acid molecule of any one of claims 29 to 31, comprising culturing the host of claim 33 under suitable conditions and isolating said polypeptide from the culture.
 - 35. A (poly)peptide encoded by a nucleic acid molecule of any one of claims 29 to 31 or as obtained by the method of claim 34.
 - 36. The (poly)peptide of claim 35 which is glycosylated, phosphorylated, amidated and/or myristylated.
 - 37. An antibody or an aptamer specifically recognizing the (poly)peptide of claim 35 or 36 or a fragment or epitope thereof.
 - 38. The antibody of claim 37, which is a monoclonal antibody.
 - 39. A transgenic non-human mammal whose somatic and germ cells comprise at least one gene encoding a functional polypeptide selected from the group

consisting of:

- (a) the polypeptide of claim 35 or 36;
- (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and
- (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution.

SEQUENCE LISTING

<110> BRAIN AG

<120> Isolation and Cloning of DNA from Uncultivated Organisms

<130> G1184 PCT

<150> 02 01 8210.1

<151> 2002-08-20

<160> 35

<170> PatentIn version 3.1

<210> 1

<211> 33925

<212> DNA

<213> Crenarchaeote

<400> 1

cacagccttg ttgatcataa cattaaaact taattcttcc agagaaattc tttttttctc 60

caagttttct gcaattgatt gcaactatctt ttttatcttt tcctttgctc tttcaaagtc 120

tttttctgaa aatatttctt taaggacatt taatatgtca tagaaagctt gtcttattat 180

tggagggtgtā tgāgācttct tccctgttaa tcctttaaca tctacagttc cgtcctccaa 240

tacacctagg taattttttt ttagttcact aaaaaccacg tagcgatata ttttatctat 300

ctccaaatct atgcctagtt cttttttaga ccaggatgaa attccactta atccttcctt 360

ggaaggattc tttaggaaca gagaatccgt atcaccgtaa ataacctcaa tcttttcttc 420

gttgcatctt tcaatagttt ttgttgttgt catccttcca accgctgcgg tagcctcagc 480

tacaggtaaa caatagagcg gaaatatttc agcaccata acccatacag tagcatttaa 540

aataaccttt atggcctgac tgataacact gtatagctgt ttatcctctt tatccaaaga 600

attatccttt gatagatatt tgtaataatt aacccttaga tcccttaggg ttcctatcaa 660

tatggaggtc atcccttgct tttccttgca aaccaatgg tttgtttgct caatatgtgt 720

tgatggatcc cttctgcaat tttcatgagg acaattgact gtttcgtaag ataaattgtg 780

aactttaatt atgctaggat acagactagc aaaatctaca actatgacat tgaaatgaat 840

tcctaaaacg ggctcaacca ccagacctcc tcgatatttt ttttccttta taatggcaac 900

tgtagacgat gttccttttt tctgtaattc atctttacgg ggaataatga tattttgctg 960

cctatgttca aaaaacatca tggacctaat ccattgattc accccgaatc ttgttatatc 1020

ttctattgac attcgggata tcctagaaat gataatcaac aattttatca gtaaattgtc 1080
attgaaagat gtcagacgaa atgtcaagtc tgcattctttg aggcaatact cggccagttt 1140
ttccaatgga agatcaccta tgctttcatc aaagtctatt tttgactcgt ttaatagggc 1200
ttcgcagata gcattttaag taaactcaga gtatttatga ctaaaagcat aattctgtac 1260
agattttattt tgaaatgtcc tgaataaatc gatatggatt ccatgcttta aggaaaccgg 1320
atccgcctga ataccctttt ttataaaaga gtctttttta actaaaatag gcaccaattc 1380
tttactaatg ggttttttgt gtacagggtc tatcgatggg tcttgagatc tagcatataa 1440
ataaggtaaa tcaaaatcat caccattaaa gggttaaaact attggataat tttgaataat 1500
agcaaaaaact tttagtatca tgtctttttc gctatcacat aattcaacgg ttgttgaatc 1560
tagtttagag ggatcaaaat ttggatcttt tcttaagacg aatacctttc taaatccatc 1620
cgatgccgat aaaccactg cagtaattac tttatcgtga tctctggctg tgggcatcct 1680
tccctcttca gagtccactt ctatatcaa agaaattctt ttgatatccg gaataggctg 1740
gtttaacaat cttgaccact ttattagaaa ctcattatat tcaactactt ttggctcatt 1800
ttctttaaaa ttaggtttta tgagattatc cagatattcg tcaactttct ctggcattgg 1860
gaattcatga aaaactaaat tattgcctat tctgttataa aacgctcccg gaattagacc 1920
caaatcgaat aaataacttt catggtattt gatatcagcc toccaagaag taaccttttc 1980
cctaaaacta ctatctgttc caccaatgga aaggggggtca ggggcaatta ttttaaaaac 2040
cgatatttcc ttgtctcaa tategtccat tttttttatt ttttctagtc taaacctatg 2100
tggtcttttg ctaactattg ttttaacctg atcagaatag agttccttta caaagcaata 2160
aggttgatgt ttatttatat gattttcaat aaaagactcg ctccaaaaat atatttgaga 2220
atcttctgga ttgtaaaact tcaaaaacac tgattttttt tctcctatgt aaacggaaga 2280
aagtaacaat gatggaatat tttctggaag ttctttttga taataactat ttttctctc 2340
cagaggcaag tcggatttta tcaccatatt tttatttcat tcatcctttt aaaaattaat 2400
gacttgtttt agaaaactat caaatttgat ttacctagca ccatcaaggc tgatgtctga 2460
ctaggttatt tttagcatta ctgatttatt tttctttgag gacaaacctt cttaagaata 2520
taccattat attattcaag cgtgtcaatc gtttaactcag tattatcaat atcgacagca 2580
atgtgagaac aagagggttt ggatgtggat ggatgatcaa tggttgattt tcaaattcct 2640
gtatgaaata ttgtgagcaa tcgatatatc attagataaa agaagattta gttaggacat 2700
cttgataata tattcgtgta gattgatttg aaaaaacagc agcaataaga aaatttaata 2760
cgataaaata cgataacttc gactaactaa catcccttgg gaaaagaact taatgcccg 2820

tccccttttt tttgtttttt aaagaaaaa gaagatttat attaacaatc ctaccataag 2880
tagtagaacg cgtccaagac aaaagggcggc gtcggtgatg aagttggttt gtgccataaa 2940
gtgatttact gacctccaga tgcgctcggt acattactgg ttaacgatat aatattaaaa 3000
tagtgggata atggggattc gatcaacaca tcggaactga cttgatggat cggatctgac 3060
atgggaagat cagacagaca ttgaaaaaga ttcattataa cagcacaaaa gatatacattg 3120
caggcattgt ttgtatgtgt gtgacaatac aatacagcat atgcttgtga gttaatctaa 3180
caaataacca aataaacaaa tcagaagggt attagagttt ttctttcttt ttcgatcgta 3240
ctctctttcc ctgcttttta aaggcaggga gaaaactcaa taaacgttct tttgtgtttg 3300
ccaatggctt tcccctcttt ctcttaccgt cttctcggtat gtgaggcggt aggcgggaag 3360
gttgttggca gagaccaaag caacgcgtat ataccacata aagcaaaagt caaccgatag 3420
gtaacaaatg ggcacggtt gtgtttttt cttgtggcg ttttgctct ctcaaaaaag 3480
gcaaggcaaa acccatatgt gtgcgtttgt catctgttat gtttttcacc atcatcattt 3540
ttttttgaat ccggttgatc ctgccggacc cgactgctat cagagtggga ctaagccatg 3600
cgagtcaaca tagcaatatg tggcatacgg ctcaagtaaca cgtagtcaac atgccagggt 3660
gacgtggata acctcgggaa actgaggata aaccgcgata agtcactact tctggaatgg 3720
gtaatgactt aaatctatat ggccctgga ttggactgct gccgatcagg ctgttgggtga 3780
ggtaatggcc caccaaact gtaaccggt cgggctctga gaggaggagc ccggagatgg 3840
gcactgagac aagggccag gccctatggg gcgcagcagg cgcgaaacct ctgcaatagg 3900
cgaaagcctg acagggttac tctgagtgat ttccgttaag gagatctttt ggcacctcta 3960
aaaatggtgc agaataagggt gtgggcaagt ctggtgtcag ccgcccggt aataccagca 4020

ccccgagtgg tcgggacgtt tattgggcct aaagcatccg tagccggttc tacaagtctt 4080
ccgttaaate cacctgctta acagatgggc tgccgaagat actatggagc taggaggcgg 4140
gagaggcaag cgggtactga tgggtagggg taaaatccgt tgatccattg aagaccacca 4200
gtggcgaaag cggcttgcca gaacgcgtc gacggtgagg gatgaaagct gggggagcaa 4260
accgattag ataccgggt agtcccagct gtaaaccgat cagactcgggt gatgaattgg 4320
cttcattgcca attcagtgcc gcagggaagc cgttaagttt gccgcctggg gactacggtc 4380
gcaagactga aacttaaagg aattggcggg ggagcaccac aagggtgaa gcctgcggtt 4440
caattggagt caacgccga aatcttaccg ggggcgacag cagaatgaag gtcaagccga 4500
agactttacc agacaagctg agaggaggt catggccgtc gccagctcgt gccgtgaggt 4560

gtcctgttaa gtcaggtaac gagcgagacc cctgcctcta gttgctacca ttattctcag 4620
gagtagtgga gctaattaga gggactgccg tcgctgagac ggaggaagga gggggctacg 4680
gcaggtcagt atgccccgaa accctcgggc cacacgcggg ctgcaatggg aaggacaatg 4740
agtatcgatt ccgaaaggag gaggcaatct ctaaacctta ccacagttat gattgagggc 4800
tgaaactcgc cctcatgaat atggaatccc tagtaaccgc gtgtcactat cgcgcggtga 4860
atacgtccct gctccttgca cacaccgccc gtcgcttcat cgaagttggg tcttggcgag 4920
gtgatgccta attggtacta tcgaacctgg ggtcagcaac gagggagaag tcgtaacaag 4980
gtggccgtag gggaacctgc ggccggatca cctccttagt tatcatatct tgcaacacag 5040
aacaaaatag aaaaaagag aaaaatgggt gggaatgaag gaaaaactct acccaccgtt 5100
taatttgttt ccttgggatc ttggtcagct tggtttacia acatgaatgc tgcagagtat 5160
acatcacaca tgcaaaaaca aagcctgcag tggatatctgt gcaagtgtta taatggacat 5220
ggatagggat atgggcatgg atgtgggcaa cacaacacaa agtagtggtg ctagaccaga 5280
tccgctctgc tcagtgcgag tggaacaaat ttgctagtga cctgtctcta tctgaatgaa 5340
tgtgtctgtc tgtctggtct tttgcgtatg cgtacccgtc ccgtgccaaa gaggcggtcg 5400
gtggttaagc caacacaata accatcagag taaaaaaaaac agagtaatgc acgcacacac 5460
acatgtgcac acagaaaaag cagaaaggaa aagagaaaga aagaaaagga aggaaagaaa 5520
gaaaaaaagt gaatacggat gcaattcttt gtcactaaac tgaggagttg gagagacaaa 5580
ggatgaagggt gacatagcag gcaacaccta gacaaatata agaaggttgt tgtttgcgat 5640
gatgtcgtgt ataggatcaa accattaaaa gatataacaa tacattagat atattaaaaa 5700
tataatgatt cacgtaaaaa tgaaatagtg aaaaattata aaaaaatgta tatctggttg 5760
tagtgcagtg taatacataa aaagcgaatg attttatcat gaaaaccgaa aattaattag 5820
ttattaccat caatagaaaa caaaattgga aactggaact ggtgcaaaga cgccggttg 5880
tgatgactc ggcttgataa gcgaagaagg acgtggcaag ctgcgataag cctggggtag 5940
gtgcatgcga ccgtcgatcc cgggatgtcc gaatgaggtc tctctttaca ctoccttgct 6000
ttgttgcttg ggagagcgaa ccgtccgaag tgaagcatct gagtaggacg aggaggagaa 6060
atcaattgag attccgtcag tagcggcgag cgaaagcgga acagcccaaa ctgaatctgc 6120
cgtggttaaca cggcagagat gtggtgttgc gggtatagcg cataggatcc tgcctttgga 6180
gctgaagtgt actggaatgt accggaacag agggtgatac ccccgtaggc aaatggaggc 6240
gggattctgc tatatccaga gtagctggcc ttggcagtgg ccagtgaagg tgggtgaaag 6300
tagtatccaa ggctaaatat tcatcaagac cgatagaaaa ctagtaccgt gagggaaagt 6360

tgaaaagtac cccggaagg ggggttaaaag cgctgaaac caaccgggta cagacgtgta 6420
tggctcgaaa ggataaaatc tagagtcata cgttccgtct agaaacacgg gccagggaga 6480
ttgctgtcat ggcaagctta accttttaca aagggaatgc gaagggaaac cgaatttgcg 6540
cattttctct ttattgagaa aagaggcaat ggatctgaaa ggggtctcaag tcatggcagt 6600
aaggctagaa accggacgat ctattcctgg ataagacgaa ggtgagtga aactcgctgg 6660
aggtctgcaa gggctctgac gtgcaaactc gtccctgat ctgggattag ggggtcaaaaa 6720
ccaatctagt ccggtgatcg ctagttccca ccgaagtga tcgcagtcct gccttagctg 6780
agatggcctg tattgtagag caccgatcgg gcggaaggg ctcgaaagag ctcgccatcc 6840
attcgaactc cgaatatacg ggcgtcgtag aagctaggag gcgggtttat gtggggtaag 6900
cctcataacc gagaggggga caaccagac taaagttaag gtcccaaact gtctactaag 6960
tgtcaaacca aagggtgttt tcgagcagag acagcaggaa ggtaggctca gaagcagcca 7020
ttctttaaa agtgcgtaac agctcacctg ccgagctcga aagccccgaa aatgtacggg 7080
gctcaagtag actaccgata ctttagacca ccgacgatgt cgggtcgtgg taggtgggcg 7140
tagtgtttgg gtagaagctg ggctgtaaag tccagtggac cgaactacta gtgcagatcc 7200
tggtggtagt aacagcatag ccgggtgaga atctcggcga ccgcatgggc aagggtttcc 7260
cggcaatgcg tcatcagccg ggagttagcc ggtcctaaaa acaacctcaa cagaattgtt 7320
tgaatgggaa actggttaat attccagtgc cttgaaagt cgttaacacc tttctgtcg 7380
cttccggata gggtaagcag aaccgtcgtt ctgtccaagt attctagctt tgaggagtac 7440
cgtaatggcg agaatcaaag cgagatacga atggcccttc gcaaggaggg tttgcttgag 7500
tccaggagac actgaaagca gaaacaggga gatactttca agaccgtacc gagatccgac 7560
actggtgcc tggatgagaa gtctaaggtc tatcgggtat accgtatggc aagggaactc 7620
ggcaaaatag ctccgtacct atggtataag gtagtcctgc agtttttacg aggagtaggg 7680
attgcaggtc gcagtgacta ggggggtccg actgtttaat aaaaacacag gtggtcgcta 7740
gtccgaaagg atgtgtatg cctctgtatc ctggccagtg gcggtaccta aaacctgggt 7800
acaaccgggc taagggccgc taaacgcgag gagtaactct gactctctta aggtagccaa 7860
atgccttgtc gggtaagttc cgacgtgcat gaatggaaca acgagggcc cgtgtccct 7920
gcctacaacc cgggaagcc acataacgtg gacgaacagt ccacgaacct ctgtcgggga 7980
gagaagacc tgtggagctt tactgcagcc tgttgttgcg atatggttgc aaatgcagag 8040
agtagctggg agccgttatg gtcagttctc cgggactgat ctaggcgaca gtgtaacacc 8100
agccatttgt taccgtatcg ctaacctgct tatgcaggga catcggcagg tgggcagttc 8160

ggctggggcg gcacccctt gaaaatgtat cgagggggcc caaagattgg ctcaggcggg 8220
acagaactcc gccggtgagg gcaaagccaa aagccagtct gactggattc ccaatgatac 8280
gggattcaga ggcgaaagcc gggcttagcg atccatcatg tcctcactat tgggggctgg 8340
tggtgacaga aaagttaccc tagggataac aggctcgtcg cgggcgagag ctcccatcga 8400
ccccgcggtt tggtacctcg atgtcggctc ttcccatcct ggttctgcag caggagccaa 8460
gggtggggct gctcgcccat taaaggggaa cgtgagctgg gtttagaccg tcgtgagaca 8520
ggtcggctctc tgcctgacag gggcgtgggt gtctgagggg aagttgcccc tagtacgaga 8580
ggaacagggc agcgcagcct ctggtttatc agttgtccga cagggaagc tgagcagcta 8640
agctgttttag gataactcct gaaagcatct aaggaggaag ctttcccg gacaagacaa 8700
ccttccgtaa ggagaagggc ggccatagaa gatggcgttg atggaatgga ggtgtaagca 8760
ccaagcttcc aagcgagggtg ttcagcctgc catcaccaat agcccaacgc acctgttgac 8820
aaaacaaaaa aaaccgacag acagaaaaaa ttgaaaatct ataactaaat atacatattt 8880
ttttgttgggt tcattatttc atgcgtaaag agtcaattat agaccaattt gatatatcta 8940
ctgattattg ttatatagaa ttttttaatg gatattgatc ataaaatttt agtatatttc 9000
atattatcta ttaacaaaat aattattaca atgggtttgg tttcggatag acaaagaaac 9060
gagacaatgg attttataaa aatactggga tataacatca gatatataaa aatagatcaa 9120
gtcaagtcaa atgaaaccat aattctgctt catggtatag gagcttccgc agaacgatgg 9180
tcagaattag tcccattttt gtataattgc aatataatta taccagacat cattggtttt 9240
ggttacagtg aaaaaccaag gatagagtac aacatagatt ttttgtaaa gtttttggt 9300
gaattgtttc tgaaacttga aatcaaaaac ccataataa tgggttcgtc ttttggtggt 9360
caattgattt tagaatatta tttcaggcac aaagactttt ttaaaaaaat gattctagt 9420
tccccggccg gtaccaaga gagaccgaca ctagcgtaa ggcaatacac ttactcatgt 9480
ttatacccaa caagagaaaa taccgaaaga gcatttaaga tgatgtcgca tttcaatcac 9540
acagtaaaag attcaatgat aaaggatttt attaatagaa tgaagcagcc caacgaaaa 9600
cactcgtttg tttcaacact tttagcacta aggaaaaata gtgatttaca agacaacctg 9660
agggaaatca aatcccaac tttagtaata tggggaaaag aggacaacac cattccagta 9720
gaaaatatag agtatttcag gggcatccct tttgtaaaaa catgcataat gagtgattgc 9780
ggtcatgtgc cttttgttga aaagcctctt gagttttata aaatagtcaa agagtttctc 9840
gactcctaatt ttctaataata agtattatat tcaacattaa aatattattg aatcaatcca 9900
cttctatgag taatgagaat gaagaaaata aagatataga ttttaagaaa tccattgaaa 9960

aggctgcgga attccagcag gatttggtgc gacagttctc tacaattcaa tacaatgcgt 10020
ttcagaatat gttttcatct ttgcaaggat ttacaaatta taatgccatg tttaaaacca 10080
ccgtacagac ggggtggcagg atctcaattc ccgaagcaga aagaaatgct ttggggattg 10140
aagaggggtga tctagtccag gttataatta taccgttgac aaggaaaaag aaaaacacaa 10200
gttaaaataa caaatccgtt aatgtgtttg aatccatttt ccaatttttg gtaaaacatt 10260
tttctgtgaa aagttgctag caattagccc tacatgccct gttggaaatt tcatcaggct 10320
tttatcctga cttgaaatta gggtgttttag ggagctactg ctgtcagacg ttacaagggtg 10380
gtcāaattca gctacaacat taagaacggg aaccttaatg ttgacaaat ttatcttggt 10440
ttcaccacaca atcatcttgt tttttgcaaa aaggttttgc tgatagatat cctttaccca 10500
ttgcctaaag gtttccccg caataggagg tgtgtcatac agccatttct ctattcttaa 10560
aaagtctgtg aaaaaacttt catcttcaaa gtttttaa ataatatagt atttgtttac 10620
accttgcttg aatggtttta gtgatgcata aaccagatac agtaattcat atggaaggtt 10680
ttcgtgatag gacagtactt tgtcaatata catgtgtcga gccatgtttt ttattacgga 10740
tttgtctttc tcggcatcaa caattggagc aatggtgact agatttttaa tgtttttttg 10800
atatagcgaa gtgtacatca aggacattgt acccccatg caatatacctt gtaatgaaat 10860
ctgatcaatg ttttctatgt tttttatgta ttctacacac tcataaataa acaaattgac 10920
ataatcatca acagtgatgt atttatccag ttttgacggg ggtttccagt caatcagata 10980
gacatttatg ccctgctcta gcaggttcct tatccaactt ttgtcgttct gcagatccaa 11040
aatatatgat ttgtttatta atgcataaac aatcaacaaa ggtacttga aagtttggtg 11100
ctttaggggt ttataatgta gtaaaccgaa gaggggtgtt tcccttatta cctcatatc 11160
gcttgatcca gtctttatgt tttctatatt cgacaatttt ttctgattt cttttaattt 11220
tgaaatgta tcaggatccc ttacaaacgt aaaataatca ttactaaat aattaattaa 11280
ggaattatc attttttttc tcatttaatt ttttttgat ttccaatgat atttttttga 11340
tttcatagag attataaaat aacaggctct tttcttctt tgacagttgt tgtgtgacc 11400
taaacaaaac ggcatttgaa tcgtaaattt ttgataact ttgatgaca tcaatgctgg 11460
aattcaataa attgttataa ttgattgaaa agtctgttga ctgcaacatt gacgagaaca 11520
catcctcaaa agtatattatg attattttcc taatatcgtc ggggtttttt tcatctatag 11580
cagacgatac cttgttttac gccacaaat aagcatttat catacgggaa aagtataggt 11640
tgagaaatga ctgatatctc agtaacaaat taccgtgttc ctttaatgta tttaaattaa 11700

ggtttggatc attcattaaa gaggtgaatg gcccgaaggt tgtaaggcta ttttaactctg 11760
ttaattgttt tataaagttt tcaaatacag actgaagtcc tgcctcttca gatagagtgt 11820
ttttactgtt tttttctctt ccaatattgt tattttctaa ttgcaaatga ttgattcacc 11880
tcggattcga aactcttgta attgtaataa tttttcatca acaatactat attctttact 11940
ttttgtcatt ataaactttt tttagggtta ggaagaaatg aaatttattg tcatttaaca 12000
aacttgaaaa gcaattatcg cagggtcaagt gtatgttaca ttgacaattc ctttttagtag 12060
agagcagttc ctaaaaactg tcaaaatgga cggatggtta aaaccattt ccaggcaaaa 12120
gcgcaactgg aatcagaaag tgtttattcc aaagtttgac catccctctg gactctattt 12180
taacgggtct gataatttta atacccttcg gttatacaaa atatggcaca tgacactgat 12240
ttaatgcttt tatgctttga ttatcacctt tttatattga aaaaacgact catgcattaa 12300
aatagtaaaa ttattctaaa taattttaga tattcacatt taggcaaata attagaagat 12360
aaagccaagg caggcccaac aattttataa ggtaatccta ccaagacgac tgggggtccg 12420
tagctcagca tggatagagc gcctgccttc tatctttacg ggagatagcc ggaagtcgag 12480
ggatcgaagc cctccggggc cgttaacctt atgggggttca aatatttttt cagttctaatt 12540
gcccatagaa ggagtatgaa tggacacatc attgcctgaa gttaacaaca accatcacia 12600
aaatgaagaa gagaaaggca taattagtgt acgaatgtga tgtcggagtt cttttattgt 12660
ttggccgggtg tctacgcgtc agaaaactgc caaaacgggc tgctcgaggt actggcaaga 12720
catcccgaac agccttgtga tcttctttag ccaactgaaa acatgatggc gtaaggggct 12780
acctcccttc agcgatatat cgctaaccgc actgcacttc cctagcggtc agatttcaag 12840
tacacagcca tcagtagcct ctaccctgta cgcgaaatgac gatattcctt gttgcaaact 12900
tttgttctga atgacatccg tcgcagtatg cagcactagc caggcgtatg catccttcag 12960
atcgccaaag caaaagtgga tgtattcgtt accccttcca tgtgagcaga cgtccctcac 13020
attgtcactt gcactcctgg gcgtggggcg ctagatttgg gttgoccagt aatcattgag 13080
ggcccaaagg cgggggtttgt ttctgggtgg cccggcataa tttgcaaaaa aaggagaaaa 13140
agaacctgac ccttgtcaaa aaatttatat ataggtagag atggttgtct atttttgtat 13200
acaggcacat gagtaattac ccaaataatta tctatggcga aaccatatat agaaacaatg 13260
agccaaggca gacatttgcc aaaaccattc gtttgtggat ggattctttc tattgtcccc 13320
cacataaccc tcttcccgag gctgatacag catcttagtt tttgggtatg cgcatacccg 13380
cccaccatgt caaaagaatg tatacatacc accgctgggt tggacatgta taaaacatct 13440
gaacttgtag gcatgtctca tgcaatgcgc catatggacg atgcggtgac ggtggatttg 13500

ccttactgct gcaaaaactc aaacaaggta ggccccattg gaaggacagc atgttctaag 13560
tcggggaaaa cggggcatgc gggtcatttt caggaggaga gtaatatccg tagcccatg 13620
ccaaatgctt gcagtctcaa ggccttgatt tgcgctctgc gcgtttgttt ctttttgaat 13680
acttaactta aggtctgggtg cctgttacag ttgatattcc accataggtt gtagatgcag 13740
agttattgct attatcagaa gtcggatttc cgggtgtctt gaaatcttgt atggagtgag 13800
tgtgctttgc ggtaccatct atcttaacca tataaaaagc agcattgaag acaactgtgg 13860
tggaattaat atttgcgaat ttatatacgc caacaactat ccacgcagga tgagttgaac 13920
taccattggt tgttgtgctg gttatagccc tatagccctt tttatatcat tctatcaagg 13980
tttgtaggat tttgtgaatg gcatatttga atctggctca tcaatatcac aatggcagaa 14040
attattacta ttgaagacaa tgataggagt ttatgtttat tgccatgcat attaagaact 14100
gaattaaaaa catggcatat aaaattctat ttttgtgact gggataaaga ataacatatt 14160
ttgtgttatg ggtcactact aagaattgag actcatcata gttgggtgtt ttggtttttag 14220
aatcaaaaaga agaaaagaac aacagcaata caagtcaaa aagacgttac ccgatcgcta 14280
atgacaatga tttggaactt ctttctattt ttaactattt tgattaggtt acagattcta 14340
ttgttgtcat tacatTTTTT gccactgtct gccagtttgg gagttggaaa ctgacctttt 14400
ttccttcagt cgcctttttg attatgccat atttatttag catttttacg cactcttctg 14460
caaacagggg ggtttctcca tattctatta gttttatgtt tgtattacca ttttttaagt 14520
atagttcttc aaaaacgggg agtttcagg caacagtggg aacacacgaa actaaggcct 14580
cagccacagc aatgccaaaa ccctctctgg atgatggaaa aataaagact ttagatttag 14640
aataaaggct aatcttttct tcctcggaga caaagcctct gtgatctata ccgcattac 14700
gtagttttgc agccttatca gggggtatgc gcccaaccat tacaaaatta gattctggtc 14760
tgagtgtttt tattgcagtc caaatttcct ccagtcctat aaatttttct atccttccga 14820
tacaaagaaa atcaatgtcc tttttattgt tgattactcc tctgttgga tcttttaaaa 14880
agatattttt atctattcca gttcctacaa tggcaattct gttggtcaga ttttttgcta 14940
attccctggg ttttttattt gctgtttcct tcaaattatt gattttgcta acccctattc 15000
catagactgt gttgagttca tgctttgacg cttgactcac ggtcaaaatc atatcagaat 15060
ccttaagcat caccgcagta gccttttgta ttagataatt gtacagaaat tcaaaatagt 15120
tcttgcaaat agatatccgc ggttcattgt gatgaaacac aacaaagatc ttggtttttg 15180
gcttgaatag cctaagtagc agccaaagga caatattaga ttcgcccga gaatccaaaa 15240
tggcaatata tggacgtttg gcaaggggcc gttaaagcat tgtatgtgtc ggttataatc 15300

ctatttggtta ttttttttgc ggagttatgg ttgtatatct tcaagacaga atagcagccg 15360
cttttgtcta tcaggtctgc aacttttttc atccagagaa atcctccggt gtaattcttt 15420
aaacttgaag ggtaaccaa gaataataat ttaatttggt ttgcgttgct tttaatcatc 15480
ctcgttggtta tatataacta taaagaacac ttattttata attgttattg ttaagaacaa 15540

gataatcaac attttggaca aacatgataa aagaattgaa aaaatgacac aataaattaa 15600
aatatttggga taatcgtcat ttacccttca aactgtgga cgatgcaaaa tatcgttttt 15660
gcagtattca tatttttctt tgcgcatatc catgaaaatg aggtttggag gttcatcagt 15720
tccttggcaa tatgtttttt gaaatattct ccaaatacat ctgtatatgc ggctccaaac 15780
tccagatttg ttccttttca aaaagatgcc catatctttg tgagctgccg ataagctccc 15840
tttttttcaa tttgtccaga taactagaat taaccttgga ttccgtaaaa ccatttttca 15900
tggccaaagt attgaggggtg ttgtgtatcc cagaaccatg ctgagctgct tcctttattc 15960
tatacgtat ctcttttggga atccctagtt tttctgcaag tttgcggtga accctttttc 16020
ctaggttgtc atagtattg ccattgtttt gaattattgag tcgcggtatc attctcagta 16080
ccgtgtctat cagattagta tctaaaagg gttcgcgtaa ctctatgctt tgagacatgg 16140
ttattttgtc ctctctttcc agtgtttctt tgtaaagtaa cttaatgtcc tctatcaggt 16200
atccctgaat ttttctgtat ccgtgttttt taacaatttt ggaataccag gaatatccgc 16260
caaacagttc gtctgcccc tgacctgtaa gcattaccog tattccctgt tcgtgagcca 16320
atttaaccgc gccatatatt ggaatggcaa cctcaacctg tccatgttg tcatcttcaa 16380
ttatgctgat tatttttggga atggtacttt caacatcact ttcagtcac tgttctattt 16440
ccaacttgag gtcaagtttt tctgctatct caagttagtt gaggatatca ottgaacctt 16500
taatcccaga cgtatagcaa ataactcgg gggccatttg ttttgccaaa tacgctacaa 16560
ttacactgtc aatcccaccg gagaaaacaa taccgatttt tttaaagtca ctcacacgtt 16620
ttctcataga ttcaaccaat gtatcaccat atgcgttaac cgcagaatcg atgtctgtgt 16680
acaggattga atattttctca catattgatt tttttgtatt tacagaaatc ggaaacaatg 16740
tagtcttgaa attggaggac ccttccttcc gcgaaatgac aagagcatag cctggcaaaa 16800
gtcttttgat ttggtcggac atagcaattt tccataaggc ttttctttct gatgcaaatg 16860
caatgaaatc actactttca ccatagtaaa tttgtcttac tocaatgcc a tcccgtacca 16920
gcacaatatc tcctgtggac tgctctctaa tcgccaaaac ataaattcca tcaagctggg 16980
taacggttct ccttatagct tcgattagat ogcctttagt gttttgataa tggctcttcaa 17040
gaagggtgaac aataacttca ctatcagtcg aggtagtaaa agtgtgatgt gcagaaaggt 17100

tctttctgat ttctttatag ttatatat ttt caccattatg ctccagaatg agttttttat 17160
cacaactcac aaacggtgc tgaccacagg agccaccaac tattgccaaa cgactgtgac 17220
ctaaaacgtc atgcccctct acctgtgaaa acaatggatt atcaaaggta tcagaataaaa 17280
ctatttgatt ctctgtagac aaacccatgc catccggacc cgggtttttc atacaggata 17340
gcatttttcc tatcaagggg gcaacatttc tctctttttt acttaaaatt ccaacaattc 17400
cacacatctt aaaattttcc tatacgggat ttattgatga acaaaatata aaagtaacca 17460
ctattgttgc cattatgggt tctaaacggt tctactctat aaaatcaagg acaccaatca 17520
tatccggtgt gtttacctta tttattgttt catttatata tttatcctta ggcatatatg 17580
cgatacctat tctgctgc tttatcatgc acaggtcacc tttagtatcg ccaatagcaa 17640
ttgtattttt tatgtctgca cagattttct ttgcatggat ttccatgtga tatctcttac 17700
acacagaatt cttgcaaaaa cagtctat tttcccatcc taacggcata tttatttctc 17760
cggtgactat ccattgtct acctcaatt catttgata aaaaaagtcc aaatcaagtt 17820
tggtcaccaa ggcctgagca gcaacactgt aactatctgt aattatccct attctgaacc 17880
cttttttctt cagcaaagat atcacctoct ggctgttctt tgcagggggg atggagtcca 17940
aagcaatttc tatttccctt tcttctattc ccctaatac agcggctatc ttctgtgtct 18000
taacatagcc tggaatggat ttgtcggact ggatgtgtct gacctgagca tacaagccaa 18060
acttttttga caatacctca attagccttc catcaattag cgtcccatcc atatcaaaaa 18120
cggccaatgt agatttaaat tccataggat acaacaaaca aggaatgtca aagaatatta 18180
ctatttagcg acagcctatt agccaaaatg tttttatagg ttggggacat cattattcaa 18240
ttgggatgtc ttgggcacca atttttttat ttcattagat atagccctat aaaaagggtta 18300
cattaaaaag tggtcgttag atcaatttta tgtatgtcat ttataaacga atatgcacat 18360
atagaaatat aaacacatga gattagatta tccacctaac tataccgaga ggataggagc 18420
agttagtatc catgcgcttc aaaagattta tgagatcgat tccggaaaga tgcccaagtt 18480
taatggcctg catcagcatc agtctataaa ggccttttgt tatgacgaac tgtcaagcat 18540
attccaagaa cttgccatag tcattccagt aaagaacgaa aaaatcagcc ttcttgaagg 18600
agtattgagc ggtattccaa atgaatgtct catcatcata gtttccaata gccaaaggac 18660
tctgtogac agatttgcca tggaggttga aatggttaagg cagtactcta gttttgcaga 18720
caagaaaata atgattattc accaaaatga tctgtagctg gctaatactt ttaagaaaat 18780
aaagtataga tccatcctca acacaaaaag tcaggttcgt agtggaagg ctgaaggaat 18840
gataattgga atattgctgg caaaaatgca cctaaaagag tacattggat ttattgacag 18900

tgataattat tttccaggag cagtaaata atagtcaag atctttgcag cgggatttgg 18960
aatggcaacc accccataca gcaatatcag aatatcgtgg cgttccaaac ccaaaatcgt 19020
aaacaactca ctacaattcc caagatgggg tagaatttca gaatccagta acaaatacct 19080
gaacgctcta atatcccaca tcacagggtt tgaaaggag attatcacga ctggaaatgc 19140
aggtgagcat gcattatcca tgtcccttgc agaaaatctc aactattcaa gcggatattc 19200
ggttgagccc tatgagtta tcaacatttt agaaaagttt ggaggtctac tcccatcaaa 19260
caatcctgac atcatagaaa aggttatcga aatatttcaa atagagacca ggaatccaca 19320
ctttcatgag gaaaaaggaa atgatactt ggcaggcatg atgcaagaat ctcttctcgc 19380

aataaacaac agcaaaattt gcaaacacaga actgaccagg gaaataaatg accatttact 19440
catgcttcag gtaaaacaca ataattgat gaccaaactc aactttaaga aaaaacacct 19500
tataatggat ccataaaaa taatacccat cgacaaattc gccgaatttg tagttaagaa 19560
ttctaaaacc ttcattagaa ttggataaaa atatgcagga atgcatattt ttgagaacaa 19620
caggtttggg aaattttgac tgatttttta gatccctcaa actgcacctt tatccatcct 19680
gttttatcaa gcctgaccaa gcgaatgcat aattatccga ccgtgttttg agcaaccaca 19740
gaggccactt tttttagaaa caacgtaaag ggataaaaaa cagttgttca ccaacatttc 19800
actagctggt gaataaatta tatcttcaaa cctttattct ccaccctac aaaccgaagg 19860
atcacagtac tcgccatcg ctacctgaaa aaaaataagc aatagtcagt ttcggaattc 19920
aaaatttcaa attttccaga gaattaattt tcccctcatc atcaatgccg tcaattactc 19980
tgaagggtat ttttccagct ctttggtatt ttgtttatt acatattttt ctggatcata 20040
tccatatttt ttgcctaact cattcatgta ctctgaaca tcatgccaag ctttttgctt 20100
gtcatatgga ttgccattta ggctgctacc tccactgcc aacgtagatt ttgatgaagc 20160
atccaatgca gtttggtata gctttgatat cttgctaaat tctcatcag tcaaccttat 20220
atttttatta ttgttggttac tggctattta ttattacccc attagtaaatt atttgatggt 20280
caaacttatc tttttctttt gataaaatgg agtcagcctt ttagcacat tttggatatt 20340
aaaccaata cgacgcgtta cggaaaagat aaaagcacct aacacccttc aaaaacattc 20400
aacgatatga ctgaaagtag ccaaagaatt tgagaatatg ttctttctca tttatcagag 20460
actttttggt tgggtttata attaattgat taacgttctg attgataaaa aagcgcaaaa 20520
tagcaaacca tgtaaatttg aaaaggggag tacatttggt tatggcttaa caatactgtg 20580
gttgtctcca aaatagtaaa ttttataatc taaaagtaga aaattcccta tgagtgatgc 20640

tatcgaaaat gtcctgatcc ttcagggagg aggatctttg ggtgcatttg gttgcggggg 20700
ctacaaagca ctagtaaaca ataacataaa acttgatatc ctgtctggca catcaattgg 20760
cggtttgaat gccacagtta ttgccggcag taaagaagat cgtccagaaa aatcattgga 20820
gaatttttgg atggaaatag ctgatactaa taatggtaat. attaatatcat accttaattt 20880
cccccttttt gaaagtcctt ttcctgggca aattcctttc cccttggcat cagaatcaac 20940
actatcattc tacagctctg ccatttatgg aaatagaaaa atctttctgc caagatgggg 21000
acctgaaaat atctttaaag atccacagta tttcacacct agcaaagga catatttgta 21060
tgaccattca cctttggtta aaaccttga aaagtacatt gattatagca aattacagcc 21120
aaacggtaag cccaacgcaa ggctaataat aaccgcagtt aacgtgatga cggcggagcc 21180
ccttattttt gacagtgcc aagcaaaaat aacccccaaa cacatacttg caaccactgc 21240
ctatccaaca tatttttttc aatgggtgga attggaaaaa gggctttttg cctgggatgg 21300
aagtttacta agcaatacco cgctaagaga agtaatagac gcatcgcccg caaaggacaa 21360
aagaatcttt cttgtcgaga actatcctaa aaatattgaa aagcttccgt caaacctaca 21420
ggaagtcaag catagggcaa gagacataat gttcagcgac aagaccgtcc acagtataca 21480
catgtccaaa gcaattacco ttcaacttaa gcttattgat gatctgtata aaatgctaga 21540
gtattacttt aattcagaaa aaatcgagga aaaggagaag ttgaaaaaa ttcgtgcgag 21600
atacaaaaaa gtttcagaag aacacggcgc agagattaaa ggtgtctact atataacacg 21660
ggacgagcca tccccctccc tttatgagaa tgcagacttt tcaaaaaatg caataaaggc 21720
atcgattaat gatggagaac aaaaggctga caggataata aaagaaatcc aaacgaaagg 21780
aaaacgaaaa taatgagcca gaaaacacca accaagttgc aatttcaaca accatttttt 21840
ttatttggcc tgtattcccc tttttgtcaa aatttttttg caggccaaaa tccaaaccaa 21900
aaggaaaatt cctaattgtc gcaaatttta ttgaaagtg aattatccat attaccatag 21960
agaggcaaat ccagttcgcc ataaaatcct aaacaaaaca atactttttg atccctgcca 22020
gaaaagcaac atcagctatc tcagatgtat tggactgagc gctgccatac cacgcgcaac 22080
tttgaataca tcgccacacc cataactctt ataccaaatt ttttaccac agaaataaca 22140
catgaattaa aacccaagaa atcagcaata taccattttt gcaaagtcaa ggcatttttag 22200
gtaatcgta atacaacaa taacagatta tacagtaaga tcattttggc aggcctaaaa 22260
aaagaccgtt ttatttaaga aaaaagcaat tctogttatg tgggtattat cattgaogat 22320
taaccaaatt aaatcaggtc aaatcaggtc aatgactttg cctatttgat aagtgataa 22380
taccttaggc caaacaatca gcaatatcga ttgttttttg catttaattat ctatttttta 22440

tattttctttt aaaaaacgaa tagaaataat caaatatgtc caaactaaaa tcaagaatta 22500
gaaaaatccg tgacagttat ataatatgtt aataataatc aactaatgac aagttcaagt 22560
gaaaatagta gtgataaaga atttgaagag ggcgcagcag gcacaaataa agatagaaaa 22620
agtgatccat tgaaagagta tgaaagtaaa gagccaatga caccagcaaa aataaatgaa 22680
ggagaaccaa cggctgtaaa gagagacca tcagaccaa agataacagg agaagggtcaa 22740
acaggagcag ataccgaaca agcagatgaa caattgcgta aacgtggcat gacccaaatc 22800
gattctgatt cttctaacac atctcaataa tcaaaaccaa aataaaaaga ataggatact 22860
tcatatcaat atctttattt tttcttttgt gaggcctttc acccggcaca aattcattat 22920
aaactaatcc aaatgcttcg tttcttgatt gtccaatagc aataaacagc gattagcaaa 22980
ccaaacatca acggcatcaa acatcaaaga aataaaaagg tgtgaaaaaa tacataacga 23040
cttgtcgta tacgaaaatg aaaatatgta ttaaactctc tocataacta gatggaaatc 23100
taattttacc ttcaccgata cccattgata agaatcatat ttttctttgc gcattgttat 23160
tattgtgcat agattgatag aaatttacga cgttaaatat ccaggagggt gctatatggt 23220

caatttttga atttgaaata ccatgtgctt tatataaaat atttcacgat atatttaaaa 23280
attacaaaaa aacatcgttt gctcgtcatt aaaaacgcaa aaaaacgggc aaaaaaata 23340
ttatatgcat tatatataga atattgtccc tatagtgttt acaatgatac ataacttta 23400
attaacaaca acccttattg ctttgcttat tgttccaata attccaatga tgacctggg 23460
aataattcca gatgtgattg cacaacagaa caogacagga attgcagact tgactgaaag 23520
caatggcggt ccagatgctg ccgttggcgg cagtagtggc accaacagta gcatagggtg 23580
taacactagt ggctcaagtg aaactatgag tggtataaac ggtggcgaag gcaccgtaga 23640
caaatttcaa tgaggcattt tgccaccct caactttatt aacggccaac cttgccgtct 23700
ataagtgatt tgtgacacct tttcctttt attattgtta ccttagttga tatcgaaaag 23760
agactgtatc caaccattta aaatatttgg ttattatggc gggttcaggc ctaaaaaag 23820
gaactgcagg caaatagggt tggcctgatg ccactttttt tggtttgct caataccaca 23880
tctattattt caactccaca aaaacattta cagggatgcc atttgctctc agtccaatcc 23940
taatccttcc ctgctcatat tcaatgtccc caatactatc ttttaatccc cttgaagcaa 24000
aatgtaaacg attacttttt agctttacct tatatgttcc cgttttttca ttatgggctg 24060
cagacacaaa ctgtatattt tctccgtcaa ccaactcata aagctoccta caaacagggt 24120
tgattacctc ccatgtttct ttatctgtta aatccttgtc caataccata aacgggtgct 24180
tcctattttt attgtttaga tatatttgat tatgatatgc agtaaagtat atttctttga 24240

caaattttaa atcgatttgc tctatttggg taaattataa aatcatacca acaataagga 24300
ttttcacaga tgatttgatt ttcacggtga atctgcccag agggatttca aaaggattgc 24360
cagcttatag ttttgtttaa acatttatgt atttaagtag gcataatcag ttagttactg 24420
tggtttcata cacctaaatt tttgccatat tgtattaaca aaattacggt aaaaaaatac 24480
cacgcaataa atcacttaaa atgtaatacc tactttccta tattgccttt ttcaaaacgc 24540
atttaactcg ttattgagaa ctattaatgt tgaatccaaa tggaacttaa cgcagcagta 24600
attgtgaaac tcgagccgga tttttctgaa gggaatgtaa gotataattc agaçggaaca 24660
cttaacagag cagaaacaaa aaacattttg gggccccata gcgcagcagc atccctagca 24720
gccctgtact caaaagtaaa acatggaacg catgtttctg tgggcacaat gggctctcca 24780
atagcagaat cgcccttaca gcaatctcaa ctgatttgcg acgctgatga actgcatctt 24840
tatagtgatc gcctctttgc aggagccgac accctggcca cagctgaagt tttgatagca 24900
ggaataaaaa aaatggcaaa tggccaagat gtggacattg ttttctcagg gcacagggca 24960
tctgatggcg aaacagggca aacaggaccc cagacagcat ggaaattagg ttatccgttc 25020
cttggaatg ttattgatta cgatattgac gttgtgaaga gaattgtaag ggtacaacgt 25080
ctaatacaaga ttacggtca tctgatatt atagaggaga tggaggcgcc tctaccggtt 25140
tttatcacac tggacccatc ctacaatccg tcttttaaca cggatatcca aaggctcaga 25200
ctagcacgaa acctacagga agcccatgat agatcacaaa ggtataagga atatctcaaa 25260
actttcaatg ccatggaact agaagtcaat ccaaagtctg tcggactgcc tggtctctcc 25320
accatagttt ataaagttga aaaaatacca agggcaaagg caaatagaaa agcagatgtt 25380
gtggatgggt ctaaccagga tagtctaagg caggttgcac gccgaatcca tgatgtttta 25440
gggggtgtag tcataaagtg acatcatcac tatctgccat acctgacgct aaactagacg 25500
aaaggccaaa ccaaatgcc catgttaatg acaaccaga aaaagaaagg ggagacaaca 25560
acaggcatct gtatgttgtg atagaacaag aggaaggcac catattacct gtgagttttg 25620
aatgcttgg tgaggcaaga aggctaattg atgattttta tcacaaatac aagccagagg 25680
aaaaagtgg tgcgattata ctggccata acatcaagca cctgtgccag gaactaatcc 25740
accatggtgc agacgcagtg atttatgccg accaccgga gctccgccac ccaagaaatc 25800
ttctttatac aaaggttgtc tgccaaattg ctacggacaa agagagcgcc gccagaattt 25860
ggccatcaaa tcccgatttt aacagacccc gttacatgtt tttttccgca gatgacacag 25920
gaaggcatth atcatcaacc gttttggcag aattgcaatc agggctggca tcagacataa 25980
acaaacttgt tatcaatgat ttagaaataa ggcatgaaca caagacaaag ggtaaaccba 26040

ttgtctatga aaagacactt gaaatgtaca gaccagactt ttcaggcttt ctttggacca 26100
ccatactctg cttggataat ataaatcccg agaacagaag gaaattccat ccacaggcat 26160
gcagtataat cccaggcgctc tttccccaaa tggaaggaga tacggataga aagggtacca 26220
taatagagtt cagcccaacc atagcccagg aagaccttag aataaaaata atcaacagaa 26280
gagtaatcaa aagcaaagtc gatttttagca ataaaaaat aatcgttagt tttggaaggg 26340
gaataaagga gtctcccgaa caaaacataa aactgataga gaaccttgca aaggaaatag 26400
aagcagaaat aggaatatca ctgcccattt caaagaaacc ctatccaata agcgaaagtc 26460
tgtcgtcaac ctatatgatt cctgacaggg ttatcggcac aagcggaaga aaggtaaadc 26520
ctcagggtga ttttgcaata ggaataagcg gggctgtcca acacatagcc gggatgaaag 26580
aatcggaatt tgtgatttcc atcaatccag acagtgaagc tcccataata gatgaatccg 26640
atgttttaat caaaggaaaa atcgagcagg tgctgcctct cctgataaat gaattaaaaa 26700
aatacaaaga gagactgcaa ataccacagg agatagaatg acaatggaaa gttttgatgt 26760
ggcgataatt ggtggagggt ctgctggact tgcggcactt gagcacctct ccaatttggg 26820
aaaacaggca atcctcatag aggcaggaaa aaaaatagga accaaaaacg tgtctggggg 26880
catattgtat tccaaaaaaa cagcaactgg aaaggccac aatgtagaag atgtgtttga 26940
taattttctg gcagacgctc cgctggaaag gaagataata aaatacatgc ttcacgccgt 27000
ctcaagggaa aaagcgttct ctctggacct gactttggca cagactatc aaacgaattt 27060

tgggtacacc gtcctgctca acaaactact ttcattggtt gcaagggaag catctcaaag 27120
tgcagaaaaa ctgggtggag ggataataac aggtgtccat ttaaggtcga taatctggaa 27180
agatgacagt accataatta tagagacaga tgaacttgag ccgttcagg taaaggcagt 27240
cattgcagct gacgggggta actcagaggt tgcgcaaata acagggtgcca gaagcaagtt 27300
cacaccgtct gacctctacc agggcgtaaa ggtggtggca aaattaccag aggggttgct 27360
tgaagagaga ttcgggggtct cggaaaacga gggagcggct cacctttttt caggcgacat 27420
aacgctaaac cacattggag gagggttcct ttacacaaac agggacacca tctcaattgg 27480
cgcagtatac cattatgact ctctaattga aaagcctaca gagcccaatg cgctgggtcaa 27540
tgcgttactg tcaaaccgtt ttgtgatgga attgataaag gacgagggtc caaggatcaa 27600
ggaggactac agggatcttt caaaggatga agaactaagg attaggttca aatccaataa 27660
attgataaaa agctggaatg acctacacca cacatattat tcaccatctg ccgttgacga 27720
gcttgtggcg cagggaataa acaaatcaag ggaggagatc aaggacaaaa ttgattcatt 27780

gtacaatgag cttgtaacaa aatacaacac agaatttgaa acaaattacg tggagttaga 27840
gtacagcgcc aaactgggtc cagatggaaa aagggtgcaga atgaaaaaac cctacttta 27900
aaacatctta ttgtcggtg atgctgcggg caggggcatt ttccttgggc cacgcataga 27960
gggcctcaac gtaggcattg atgacgcggt tagggccgca gaagctgtct caaagtcaat 28020
agatcaaaat aactttcagt ttgacaacat tggngaacgc tacactaaat cagtggatga 28080
aagtcacat accgcagaca tgagcaggat cgacgcaaac tatctcaaag ccgttcttga 28140
ttgcacaaaa aaggttccca aaaacactct tgggtttaag tatgggtcta ttgtcaaatt 28200
gatgtcaaat agcaccttta ggaatgtatc cataggaatt gcaaactcta tagggtaaca 28260
aaggctttta cctgtgattg agtcagacaa aacctacaat caaattccca tcgagattgc 28320
ggagagaaat ggcaaagatt tgcggaaaag ctattccata gagattccca ccattgccga 28380
gcgtattgct aatctgaact ataatgacga ttcactgtca cacatcaagg ttttgaactc 28440
gcaaagtgc tttatgaaaa aaatggtcca actgtgccct accaaatgct acagtattga 28500
gaatgagcgg ataatgctac agcacgaagg atgcatagag tgtgggacat gcgcaagaga 28560
aacagaatgg aggcacctc gtggggaaaa aggaataatc tataattacg ggtaagccat 28620
aaccggaatc catcaacata tcctttctgg aaaaaagtc ggggataaca cagcaacaa 28680
aaaaaacaac gaatggattt caggttctaa atttttgggt gtttacacct tatctctgct 28740
ttcaccgctt ttattttttt ttgatggat tctatttctt caatcaactc atcaacatac 28800
tccttgattt ttctgtggct ccccgataac cctccagccc ctgcatactc agttagagaa 28860
tcagtttggt tccggatcac tgtctgcatt tttcatcaa gctcttcaat caactgtatg 28920
attgacggaa taagcctctc catatgattg gtaaagccac caacaattat aaagcttacg 28980
cagtgtatta atgctatta cagtctatat gggtataaac aaccaacaa aatccgaatc 29040
aaaagtaa at gaataacaca taatactaca atgggccatg aaacaaatta catcaaagcg 29100
tcacatttta agcaacgtca actgctagtt ttgaaagtta tgtatttctt tagattattt 29160
tctattctat ttcatgtgt gtagttgggt gttgcagcag cagttgttgc agttgccaaa 29220
ctcatgggtc ttttccttca tcgttttttt ctggcatgtc ttttgcggc atatgggaag 29280
gcaggagac aggtatgaca aatttgaatg tggcgccat ttttccttca ctgccaaggt 29340
ggtgcagtat ttcacaaatg ccttcgtctt tatctttagt gttgttctg ctgctgctgt 29400
tgccgccctt gctgtttgtc tcctcaaacc atatttttcc accatgctcc tcaacaattt 29460
tccttgacag gtataggcca aggccggtc cctgggttga ctttgtgaca aatttctgaa 29520
acagctgac ccttattttg gagttgagcc caaccccggt gtccctgact gtgactagca 29580

ccgcgccttc tttctgcctc ccgatgtggt caccaccatt gtcaccacca ccgttgctgc 29640
tgtcgtgct gctatccact ctgccccat tgcctttacc agctgtagca gtgtttgagg 29700
tatcactttc ctgagaggtg gaagtgaagg gagaagactc acccatcact gccgtggaaa 29760
caacaatctt gccgtcattg gtgaacttca ttgcgttgct cagcaggttg aaaacaacct 29820
ggcttatctt ttgcgatca cagtctacat aaaaagggtg gttggggcca tttacgggtt 29880
ctaccactg ctctttttgc tgcgtctctt tttgcgctg ttttgcctc gccgtgcct 29940
ttacccttc tgcctttgcg ccgcgccac ctctggagta ccgccattt ctgttgccgt 30000
cagatggcaa aaacactatc gccacctgt ttgccttctc ctgttaggcg tttttttct 30060
caatgtctc tatcacctgg gaaatcagg tgtggatata cacatTTTTT tggatgtcca 30120
ggctaaagct tccgctttcg attctgtca cctgcagaat gctttcggca aggttctgca 30180
gccgggacgc gtttcttggt atcatgtcaa gctcccgctg aaactctgtt tttctttgc 30240
caagcttctc ctccagtatc tccacaccgt ttaggatggg catgattggc gttcgcaact 30300
catgcgcagc cacgtttatg aactcgcttt tgaactttgc gttttggtca agctgctgga 30360
acaggacact ctggtcatag aggacctcaa atatggacga gtaagacaat accgttggct 30420
cgctgttga gtagattgaa aagccgattg cggcggttgc cacctctcc ctgctgtga 30480
tcagctccat caccagcgac tcctttctgt ccacaaccag tgtctttatc ttgatgcaa 30540
tgcttgccgc aatgtcctgg acttggatgt tgggcctgta ttttgtgagg agcctcaagg 30600
acaaggactc tgcactgag gcatccatcg gcgtgaggat gttgatctc agctgtcgt 30660
tttgctccac catctcttc aagagttgca gtgtgccgc ttttctctgc aggtggaacg 30720
cgtaaacctg ggagtaact atcagtatct cccttttggc cctgcttata atttcaaact 30780
ccctttggac cgcgtccttg tagttggaga agaccagtga gacgggcatg acaaccccat 30840
cctccagctc ctttatcctg tgcctgctg gcaacgcct gcccaaaaag ctgtcaaaca 30900

caaaactgtg ctgctctgca atctcagga ggttctgaa caaaagctgg ggtattgact 30960
gtgccgcatg aagggtggcg acagccacat actccctctg gtccgccacc tcaaagtctc 31020
ccttcagccc atccaggtgc ctaatctccg aaaacgagag catctccttg acatagccga 31080
cgttgtcctt tgttatttcg gttacatacc gcagtttaag gccctgttt ttgaccgct 31140
caaccctttt ctcccttatg gcgtcaaccc ctatcatcac ggacggggcc acggagtta 31200
tgcaagagtc tatcttcaca ttggccctgt ctatgaacct caaaatggcg ttgtttgcgt 31260
tttcagggcc atagtacacc tttgtggtgg gtgcaccggc gtcaacacca ccatcattat 31320
cgttctcgtt attgttgacg tcacaaaac cgcgtttgtg cgacaaaccg gggtttgcgt 31380

caatgtatattt attttcacat acattatcat tatattgtgcc aatgtcatgg agattattat 31440
tatgtatttag agaagaacct tccaaccgtt atcacgatat cctattttgt tcaactatatt 31500
aatttgagct taaaacttta taaataccgt atatacggta acaggattat tatttctaaaa 31560
aaacacttaa aggtacttga caaaattctg aacaaaagat ccccatatt tactaaatac 31620
caagatttct gcaaactgat gtgatgtgat gtatgcatag taccaatata taggcaaacg 31680
tttttggcat tagaaaggaa tacgaataga taatcaaaga atgaaatggc cgaacacaaa 31740
ccaacaccat ggctttcaca cactgtcatt attatgtgaa actttatctg gcctctaatac 31800
tttgtcagga attaaactgt tttttattgc caattctata atgatatgct ataagcagtt 31860
agattacctt ttgatggttag tggttgttcc agtagtggtt tctccagtaa tatcatgagt 31920
ttaaagaccc ggctgatggc agcgatagaa tgcttaattg catctattga ggaaagtgtt 31980
gttgaaggta caacgctcaa tactatcaat tgaggacaac agggattgag aattgttttg 32040
acaatgataa tccattcata aaaaaattg caagataaag catatgccgc gattgttgac 32100
cccctatttt gcatgcgttc caacaaaaag ttgtcttaac ttgacagaca ttgaataaa 32160
ttaaaaagat gttgttgact ttogtttatt gattgattaa gattacgggtt ttattttacc 32220
aaggatttaa gcattacttg cctacacgaa attaaattgc gagcaggaaa acagggaatgt 32280
gtttacataa taagaatata cccctaacca agtctttttc ctatgcgatt tttttttggt 32340
tacgccaggc cgaagaatat acttttggtg acaatgattg gtaaaccctt taaccottgct 32400
tttgcgtaga ttgtcataat tgatgttcgt aaagataaaa gcaataaaaa gaaatagtca 32460
ttatgtagaa taacacattt tttttataac ccgttataat ttaattgcaa agcagtcatac 32520
tttctaaaaat aatcacaatt tgcagaatgc cgtcacttca tcttggtgca tatggtttaa 32580
ttttggatat ttctgaaagc ccaatcacaa gggttaaagg tagaacaagt cacttgatta 32640
ttaaaatata tccacatatg gataacaata caaggatgag ttcttttagca atcgagtttt 32700
ttttatccct tttttcaata acgttacttt ctaaaagaat ataccaacca gtgaaatcaa 32760
agtcataatac ctacatgac aagcatccat ttcagtacaa gatggaggat tatgcaaacc 32820
acaacaaaat tgtagactat aaaaactgct tacttttttt tcaagtatcg atgttacaaa 32880
aaaataaaaat aattaggatt cgggttccag gtttgtttta tacaggtggc tggatttccc 32940
tcacactaaa gtttttgata tccacatcat ttgcaccatc ccacctgaaa gtagcaatgg 33000
ggcctcccca ggatataatc tgatccggtt caccaccaca ctcttcacca tcatttccaa 33060
accacctga gtcagtgaat gtgtatacct tttgccattt gttcttcaga gtcgggctat 33120
ccgggtttct gtctacccat atttcagtgg tgactacggc ctcaccacca gccatttggc 33180

ggttatagat catggcttta aatccaatga atctatcaaa actagacgcc gaaggtgagg 33240
gtgtggtagt gcttgaaaac acataggaga catgccactg ctcttttgca agcctaaccc 33300
ttccatcata gaatagatct gctttatatg ctgagccctc gcatccttcg ccatcatagt 33360
gcctaccacc cctgtcatac caagcgaaat tttcagaatc atctccacta ttaaccctta 33420
caatacccggt catttccaca ttcttccaat catttggata ctgcatgtat ccttgtgttg 33480
cgagtaccga gtgatcgtaa gtctcaatat cctctggatg gtaccctgat gatgtaaaca 33540
cgttatatct gacctgatcg tcattaacgt tccaactgcc atctgggttt aggtccatgt 33600
caggtggggtt tgctcgtgga tcattgttcg ggttttgcatt attcataaac catttttctc 33660
caccacccgc cttatcgggg taaatctggg ttatcccaa ctggtctaata gtccctccgc 33720
ctccagaagg ggcaacagag aacgtccata ccttgtcggc agccaatggg acaccagtcg 33780
catccgtagc accggttggt attctggcag tgtatgtggc accaggtgtt aaatctgcag 33840
aggggtttag ggtcgcaact gtgttggttg gtgaattcat gcttacggtt gcgggcacag 33900
gtgcgcctcc gcttgttagc agtgt 33925

<210> 2

<211> 2367

<212> DNA

<213> Crenarchaeote

<400> 2

atggtgataa aatccgactt gcctctggag gagaaaaata gttattatca aaaagaactt 60
ccagaaaaata ttccatcatt gttactttct tccgtttaca taggagaaaa aaaatcagtg 120
tttttgaagt ttacaatcc agaagattct caaatatatt tttggagcga gtcttttatt 180
gaaaatcata taaataaaca tcaaccttat tgctttgtaa aggaactcta ttctgatcag 240
gttaaaacaa tagtttagcaa agagccacat aggttttagac tagaaaaaat aaaaaaatg 300
gacgatattg aggacaagga aatatcggtt tttaaaataa ttgccctga cccctttcc 360
attggtggaa cagatagtag ttttagggaa aaggttactt cttgggaggc tgatatcaaa 420
taccatgaaa gttatttatt cgatttgggt ctaattccgg gagcgtttta taacagaata 480
ggcaataatt tagtttttca tgaattccca atgccagaga aagttgacga atatctggat 540
aatctcataa aacctaat taaagaaaat gagccaaaaa gtagtgaata taatgagttt 600
ctaataaagt ggtcaagatt gttaaaccag cctattccgg atatcaaaag aatttctttg 660

gatatagaag tggactctga agaggggaagg atgccacag ccagagatca cgataaagta 720

attactgcag tgggtttatc ggcatcggat ggatttagaa aggtattcgt cttagaaaa 780

gatccaaatt ttgatccctc taaactagat tcaacaaccg ttgaattatg tgatagcgaa 840

aaagacatga tactaaaagt ttttgctatt attcaaaatt atccaatagt tttaaccttt 900

aatggtgatg attttgattt accttattta tatgctagat ctcaagacc atcgatagac 960

cctgtacaca aaaaacccat tagtaaagaa ttggtgccta ttttagttaa aaaagactct 1020

tttataaaaa ggggtattca ggcgatccg gtttccttaa agcatggaat ccatatcgat 1080

ttattcagga catttcaaaa taaatctgta cagaattatg cttttagtca taaatactct 1140

gagtttactt taaatgctat ctgcgaagcc ctattaaacg agtcaaaaat agactttgat 1200

gaaagcatag gtgatcttcc attggaaaaa ctggccgagt attgcctcaa agatgcagac 1260

ttgacatttc gtctgacatc tttcaatgac aatttactga taaaattgtt gattatcatt 1320

tctaggatat cccgaatgtc aatagaagat ataacaagat tcggggtgaa tcaatggatt 1380

aggtccatga tgttttttga acataggcag caaaatatca ttattccccg taaagatgaa 1440

ttacagaaaa aaggaacatc gtctacagtt gccattataa aggaaaaaaa atatcgagga 1500

ggtctggtgg ttgagcccgt tttaggaatt catttcaatg tcatagttgt agattttgct 1560

agtctgtatc ctagcataat taaagttcac aatttatctt acgaaacagt caattgtcct 1620

catgaaaatt gcagaaggga tccatcaaca catattgagc aaacaaacca ttgggtttgc 1680

aaggaagaac aagggatgac ctccatattg ataggaaccc taagggatct aagggttaat 1740

tattacaaat atctatcaaa ggataattct ttggataaag aggataaaca gctatacagt 1800

gttatcagtc aggcataaa ggttatttta aatgctacgt atggggttat ggggtctgaa 1860

atatttccgc tctattgttt acctgtagct gaggctaccg cagcggttg aaggatgacc 1920

acaacaaaaa ctattgaaaa atgcaacgaa gaaaagattg aggttattta cgggtgatac 1980

gattctctgt tcctaaagaa tcctccaag gaaggattaa gtggaatttc atcctggtct 2040

aaaaaagAAC taggcataga tttggagata gataaaagat atcgctacgt ggttttttagt 2100

gaactaaaaa aaaattacct aggtgtattg gaggacggaa ctgtagatgt taaaggatta 2160

acagggaaga agtctcatac acctccaata ataagacaag ctttctatga catattaaat 2220

gtccttaaaag aaatattttc agaaaaagac tttgaaagag caaaggaaaa gataaaaaaa 2280

atagtgcaat caattgcaga aaacttgag aaaaaagaa tttctctgga agaattaagt 2340

tttaatgtta tgatcaacaa ggctgtg 2367

<210> 3

<211> 789

<212> PRT

<213> Crenarchaeote

<400> 3

Met Val Ile Lys Ser Asp Leu Pro Leu Glu Glu Lys Asn Ser Tyr Tyr
1 5 10 15

Gln Lys Glu Leu Pro Glu Asn Ile Pro Ser Leu Leu Leu Ser Ser Val
20 25 30

Tyr Ile Gly Glu Lys Lys Ser Val Phe Leu Lys Phe Tyr Asn Pro Glu
35 40 45

Asp Ser Gln Ile Tyr Phe Trp Ser Glu Ser Phe Ile Glu Asn His Ile
50 55 60

Asn Lys His Gln Pro Tyr Cys Phe Val Lys Glu Leu Tyr Ser Asp Gln
65 70 75 80

Val Lys Thr Ile Val Ser Lys Glu Pro His Arg Phe Arg Leu Glu Lys
85 90 95

Ile Lys Lys Met Asp Asp Ile Glu Asp Lys Glu Ile Ser Val Phe Lys
100 105 110

Ile Ile Ala Pro Asp Pro Leu Ser Ile Gly Gly Thr Asp Ser Ser Phe
115 120 125

Arg Glu Lys Val Thr Ser Trp Glu Ala Asp Ile Lys Tyr His Glu Ser
130 135 140

Tyr Leu Phe Asp Leu Gly Leu Ile Pro Gly Ala Phe Tyr Asn Arg Ile
145 150 155 160

Gly Asn Asn Leu Val Phe His Glu Phe Pro Met Pro Glu Lys Val Asp
165 170 175

Glu Tyr Leu Asp Asn Leu Ile Lys Pro Asn Phe Lys Glu Asn Glu Pro
180 185 190

Lys Ser Ser Glu Tyr Asn Glu Phe Leu Ile Lys Trp Ser Arg Leu Leu
195 200 205

Asn Gln Pro Ile Pro Asp Ile Lys Arg Ile Ser Leu Asp Ile Glu Val
210 215 220

Asp Ser Glu Glu Gly Arg Met Pro Thr Ala Arg Asp His Asp Lys Val
225 230 235 240

Ile Thr Ala Val Gly Leu Ser Ala Ser Asp Gly Phe Arg Lys Val Phe
245 250 255

Val Leu Arg Lys Asp Pro Asn Phe Asp Pro Ser Lys Leu Asp Ser Thr
260 265 270

Thr Val Glu Leu Cys Asp Ser Glu Lys Asp Met Ile Leu Lys Val Phe
275 280 285

Ala Ile Ile Gln Asn Tyr Pro Ile Val Leu Thr Phe Asn Gly Asp Asp
290 295 300

Phe Asp Leu Pro Tyr Leu Tyr Ala Arg Ser Gln Asp Pro Ser Ile Asp
305 310 315 320

Pro Val His Lys Lys Pro Ile Ser Lys Glu Leu Val Pro Ile Leu Val
325 330 335

Lys Lys Asp Ser Phe Ile Lys Arg Gly Ile Gln Ala Asp Pro Val Ser
340 345 350

Leu Lys His Gly Ile His Ile Asp Leu Phe Arg Thr Phe Gln Asn Lys
355 360 365

Ser Val Gln Asn Tyr Ala Phe Ser His Lys Tyr Ser Glu Phe Thr Leu
370 375 380

Asn Ala Ile Cys Glu Ala Leu Leu Asn Glu Ser Lys Ile Asp Phe Asp
385 390 395 400

Glu Ser Ile Gly Asp Leu Pro Leu Glu Lys Leu Ala Glu Tyr Cys Leu
405 410 415

Lys Asp Ala Asp Leu Thr Phe Arg Leu Thr Ser Phe Asn Asp Asn Leu
420 425 430

Leu Ile Lys Leu Leu Ile Ile Ile Ser Arg Ile Ser Arg Met Ser Ile
435 440 445

Glu Asp Ile Thr Arg Phe Gly Val Asn Gln Trp Ile Arg Ser Met Met
450 455 460

Phe Phe Glu His Arg Gln Gln Asn Ile Ile Ile Pro Arg Lys Asp Glu
465 470 475 480

Leu Gln Lys Lys Gly Thr Ser Ser Thr Val Ala Ile Ile Lys Glu Lys
485 490 495

Lys Tyr Arg Gly Gly Leu Val Val Glu Pro Val Leu Gly Ile His Phe
500 505 510

Asn Val Ile Val Val Asp Phe Ala Ser Leu Tyr Pro Ser Ile Ile Lys
515 520 525

Val His Asn Leu Ser Tyr Glu Thr Val Asn Cys Pro His Glu Asn Cys
530 535 540

Arg Arg Asp Pro Ser Thr His Ile Glu Gln Thr Asn His Trp Val Cys
545 550 555 560

Lys Glu Lys Gln Gly Met Thr Ser Ile Leu Ile Gly Thr Leu Arg Asp
565 570 575

Leu Arg Val Asn Tyr Tyr Lys Tyr Leu Ser Lys Asp Asn Ser Leu Asp
580 585 590

Lys Glu Asp Lys Gln Leu Tyr Ser Val Ile Ser Gln Ala Ile Lys Val
595 600 605

Ile Leu Asn Ala Thr Tyr Gly Val Met Gly Ala Glu Ile Phe Pro Leu
610 615 620

Tyr Cys Leu Pro Val Ala Glu Ala Thr Ala Ala Val Gly Arg Met Thr
625 630 635 640

Thr Thr Lys Thr Ile Glu Lys Cys Asn Glu Glu Lys Ile Glu Val Ile
645 650 655

Tyr Gly Asp Thr Asp Ser Leu Phe Leu Lys Asn Pro Ser Lys Glu Gly
660 665 670

Leu Ser Gly Ile Ser Ser Trp Ser Lys Lys Glu Leu Gly Ile Asp Leu
675 680 685

Glu Ile Asp Lys Arg Tyr Arg Tyr Val Val Phe Ser Glu Leu Lys Lys
690 695 700

Asn Tyr Leu Gly Val Leu Glu Asp Gly Thr Val Asp Val Lys Gly Leu
705 710 715 720

Thr Gly Lys Lys Ser His Thr Pro Pro Ile Ile Arg Gln Ala Phe Tyr
725 730 735

Asp Ile Leu Asn Val Leu Lys Glu Ile Phe Ser Glu Lys Asp Phe Glu
740 745 750

Arg Ala Lys Glu Lys Ile Lys Lys Ile Val Gln Ser Ile Ala Glu Asn
755 760 765

Leu Glu Lys Lys Arg Ile Ser Leu Glu Glu Leu Ser Phe Asn Val Met
770 775 780

Ile Asn Lys Ala Val
785

<210> 4

<211> 882

<212> DNA

<213> Crenarchaeote

<400> 4

atggatattg atcataaaat tttagtatat ttcattattat ctattaacaa aataattatt 60

acaatgggtt tggtttcgga tagacaaaga aacgagacaa tggattttat aaaaatactg 120

ggatataaca tcagatatat aaaaatagat caagtcaagt caaatgaaac cataattctg 180

cttcatggta taggagcttc cgcagaacga tggtcagaat tagtcccatt tttgtataat 240

tgcaatataa ttataccaga catcattggt tttggttaca gtgaaaaacc aaggatagag 300

tacaacatag atttatttgt aaagtttttg gatgaattgt ttctgaaact tgaaatcaaa 360

aacccataa taatgggttc gtcttttggt ggtcaattga ttttagaata ttatttcagg 420

cacaaagact tttttaaaaa aatgattcta gtgtccccg cgggtaccca agagagaccg 480

aactagcgt taaggcaata cacttactca tgtttatacc caacaagaga aaataccgaa 540
 agagcattta agatgatgtc gcatttcaat cacacagtaa aagattcaat gataaaggat 600
 tttattaata gaatgaagca gcccaacgca aaacactcgt ttgtttcaac acttttagca 660
 ctaaggaaaa atagtgattt acaagacaac ctgagggaaa tcaaaatccc aacttttagta 720
 atatggggaa aagaggacaa caccattcca gtagaaaata tagagtattt caggggcatc 780
 ccttttgtaa aaacatgcat aatgagtgat tgcggtcatg tgccttttgt tgaaaagcct 840
 cttgagtttt ataaaatagt caaagagttt atcgactcct aa 882

<210> 5

<211> 293

<212> PRT

<213> Crenarchaeote

<400> 5

Met Asp Ile Asp His Lys Ile Leu Val Tyr Phe Ile Leu Ser Ile Asn
1 5 10 15

Lys Ile Ile Ile Thr Met Gly Leu Val Ser Asp Arg Gln Arg Asn Glu
20 25 30

Thr Met Asp Phe Ile Lys Ile Leu Gly Tyr Asn Ile Arg Tyr Ile Lys
35 40 45

Ile Asp Gln Val Lys Ser Asn Glu Thr Ile Ile Leu Leu His Gly Ile
50 55 60

Gly Ala Ser Ala Glu Arg Trp Ser Glu Leu Val Pro Phe Leu Tyr Asn
65 70 75 80

Cys Asn Ile Ile Ile Pro Asp Ile Ile Gly Phe Gly Tyr Ser Glu Lys
85 90 95

Pro Arg Ile Glu Tyr Asn Ile Asp Leu Phe Val Lys Phe Leu Asp Glu
100 105 110

Leu Phe Leu Lys Leu Glu Ile Lys Asn Pro Ile Ile Met Gly Ser Ser
115 120 125

Phe Gly Gly Gln Leu Ile Leu Glu Tyr Tyr Phe Arg His Lys Asp Phe
130 135 140

Phe Lys Lys Met Ile Leu Val Ser Pro Ala Gly Thr Gln Glu Arg Pro
145 150 155 160

Thr Leu Ala Leu Arg Gln Tyr Thr Tyr Ser Cys Leu Tyr Pro Thr Arg
165 170 175

Glu Asn Thr Glu Arg Ala Phe Lys Met Met Ser His Phe Asn His Thr
180 185 190

Val Lys Asp Ser Met Ile Lys Asp Phe Ile Asn Arg Met Lys Gln Pro
195 200 205

Asn Ala Lys His Ser Phe Val Ser Thr Leu Leu Ala Leu Arg Lys Asn
210 215 220

Ser Asp Leu Gln Asp Asn Leu Arg Glu Ile Lys Ile Pro Thr Leu Val
225 230 235 240

Ile Trp Gly Lys Glu Asp Asn Thr Ile Pro Val Glu Asn Ile Glu Tyr
245 250 255

Phe Arg Gly Ile Pro Phe Val Lys Thr Cys Ile Met Ser Asp Cys Gly
260 265 270

His Val Pro Phe Val Glu Lys Pro Leu Glu Phe Tyr Lys Ile Val Lys
275 280 285

Glu Phe Ile Asp Ser
290

<210> 6

<211> 318

<212> DNA

<213> Crenarchaeote

<400> 6

ttgaatcaat ccacttctat gagtaatgag aatgaagaaa ataaagatat agattttaag 60

aaatccattg aaaaggctgc ggaattccag caggatttgt tgcgacagtt ctctacaatt 120

caatacaatg cgtttcagaa tatgttttca tctttgcaag gatttacaaa ttataatgcc 180

atgttttaaaa ccaccgtaca gacgggtggc aggatctcaa ttcccgaagc agaaagaaat 240
gctttgggga ttgaagaggg tgatctagtc caggttataa ttataccggt gacaaggaaa 300
aagaaaaaca caagttaa 318

<210> 7

<211> 105

<212> PRT

<213> Crenarchaeote

<400> 7

Met Asn Gln Ser Thr Ser Met Ser Asn Glu Asn Glu Glu Asn Lys Asp
1 5 10 15

Ile Asp Phe Lys Lys Ser Ile Glu Lys Ala Ala Glu Phe Gln Gln Asp
20 25 30

Leu Leu Arg Gln Phe Ser Thr Ile Gln Tyr Asn Ala Phe Gln Asn Met
35 40 45

Phe Ser Ser Leu Gln Gly Phe Thr Asn Tyr Asn Ala Met Phe Lys Thr
50 55 60

Thr Val Gln Thr Gly Gly Arg Ile Ser Ile Pro Glu Ala Glu Arg Asn
65 70 75 80

Ala Leu Gly Ile Glu Glu Gly Asp Leu Val Gln Val Ile Ile Ile Pro
85 90 95

Leu Thr Arg Lys Lys Lys Asn Thr Ser
100 105

<210> 8

<211> 1086

<212> DNA

<213> Crenarchaeote

<400> 8

atgagaaaaa aaatgaataa ttccttaatt aattatttag tgaatgatta ttttacgttt 60

gtaagggatc ctgataacat ttcaaaatta aaagaaatca ggaaaaaatt gtcgaatata 120
 gaaaacataa agactggatc aagcgaatat gaggtaataa gggaaacaac cctcttccgt 180
 ttactacatt ataaaccctt aaagcaacaa actttcaagt accctttgtt gattgtttat 240
 gcattaataa acaaatcata tttttggat ctgcagaacg acaaaagttg gataaggaac 300
 ctgctagagc agggcataaa tgtctatctg attgactgga aacccccgtc aaaactggat 360
 aaatacatca ctgttgatga ttatgtcaat ttgtttattt atgagtgtgt agaatacata 420
 aaaaacatag aaaacattga tcagatttca ttacaaggat attgcatggg ggggtacaatg 480
 tccttgatgt aacttgcgt atatcaaaaa aacattaaaa atctagtcac cattgctcca 540
 attgttgatg ccgagaaaga caaatccgta ataaaaaaca tggctgagca catggatatt 600
 gacaaagtac tgtcctatca cgaaaacttt ccatatgaat tactgtatct ggtttatgca 660
 tcactaaaac cattcaagca aggtgtaaac aaatactata atttatttaa aaactttgaa 720
 gatgaaagtt ttgtacagaa ctttttaaga atagagaaat ggctgtatga cacacctcct 780
 attgcggggg aaacctttag gcaatgggta aaggatatct atcagcaaaa cctttttgca 840
 aaaaacaaga tgattgtggg tgaaaacaag ataaatttgt caaacattaa ggttcccggt 900
 cttaatgttg tagctgaatt tgaccacctt gtaacgtctg acagcagtag ctccctaaac 960
 aacctaattt caagtcagga taaaagcctg atgaaatttc caacagggca tgtagggcta 1020
 attgctagca acttttcaca gaaaaatgtt ttaccaaaaa ttggaaaatg gattcaaaca 1080
 cattaa 1086

<210> 9

<211> 361

<212> PRT

<213> Crenarchaeote

<400> 9

Met Arg Lys Lys Met Asn Asn Ser Leu Ile Asn Tyr Leu Val Asn Asp
 1 5 10 15

Tyr Phe Thr Phe Val Arg Asp Pro Asp Asn Ile Ser Lys Leu Lys Glu
 20 25 30

Ile Arg Lys Lys Leu Ser Asn Ile Glu Asn Ile Lys Thr Gly Ser Ser
 35 40 45

Glu Tyr Glu Val Ile Arg Glu Thr Thr Leu Phe Arg Leu Leu His Tyr
50 55 60

Lys Pro Leu Lys Gln Gln Thr Phe Lys Tyr Pro Leu Leu Ile Val Tyr
65 70 75 80

Ala Leu Ile Asn Lys Ser Tyr Ile Leu Asp Leu Gln Asn Asp Lys Ser
85 90 95

Trp Ile Arg Asn Leu Leu Glu Gln Gly Ile Asn Val Tyr Leu Ile Asp
100 105 110

Trp Lys Pro Pro Ser Lys Leu Asp Lys Tyr Ile Thr Val Asp Asp Tyr
115 120 125

Val Asn Leu Phe Ile Tyr Glu Cys Val Glu Tyr Ile Lys Asn Ile Glu
130 135 140

Asn Ile Asp Gln Ile Ser Leu Gln Gly Tyr Cys Met Gly Gly Thr Met
145 150 155 160

Ser Leu Met Tyr Thr Ser Leu Tyr Gln Lys Asn Ile Lys Asn Leu Val
165 170 175

Thr Ile Ala Pro Ile Val Asp Ala Glu Lys Asp Lys Ser Val Ile Lys
180 185 190

Asn Met Ala Glu His Met Asp Ile Asp Lys Val Leu Ser Tyr His Glu
195 200 205

Asn Phe Pro Tyr Glu Leu Leu Tyr Leu Val Tyr Ala Ser Leu Lys Pro
210 215 220

Phe Lys Gln Gly Val Asn Lys Tyr Tyr Asn Leu Phe Lys Asn Phe Glu
225 230 235 240

Asp Glu Ser Phe Val Gln Asn Phe Leu Arg Ile Glu Lys Trp Leu Tyr
245 250 255

Asp Thr Pro Pro Ile Ala Gly Glu Thr Phe Arg Gln Trp Val Lys Asp
260 265 270

Ile Tyr Gln Gln Asn Leu Phe Ala Lys Asn Lys Met Ile Val Gly Glu
275 280 285

Asn Lys Ile Asn Leu Ser Asn Ile Lys Val Pro Val Leu Asn Val Val
290 295 300

Ala Glu Phe Asp His Leu Val Thr Ser Asp Ser Ser Ser Ser Leu Asn
305 310 315 320

Asn Leu Ile Ser Ser Gln Asp Lys Ser Leu Met Lys Phe Pro Thr Gly
325 330 335

His Val Gly Leu Ile Ala Ser Asn Phe Ser Gln Lys Asn Val Leu Pro
340 345 350

Lys Ile Gly Lys Trp Ile Gln Thr His
355 360

<210> 10

<211> 582

<212> DNA

<213> Crenarchaeote

<400> 10

ttgcaattag aaaataacaa tattggagag gaaaaaaca gtaaaaacac tctatctgaa 60
gaggcaggac ttcagtctgt atttgaaaac ttataaaac aattaacaga gttaaatagc 120
cttacaacct tggggccatt cacctcttta atgaatgac caaaccttaa tttaaataca 180
ttaaaggaa acggtaat tttactgaga tatcagtcac ttctcaacct atacttttcc 240
cgtatgataa atgcttattt gttggccgta aacaaggat cgtctgctat agatgaaaaa 300
aaccgccgac atattaggaa aataatcata aatacttttg aggatgtgtt ctogtcaatg 360
ttgcagtcaa cagacttttc aatcaattat aacaatttat tgaattccag cattgatgtc 420
atcaaaagtt atcaaaaaat ttacgattca aatgccgttt tgtttaggtc acaacaacaa 480
ctgtcaaaag aagaaaaaga cctgttattt tataatctct atgaaatcaa aaaaatatca 540
ttggaaatca aaaaaaatt aaatgagaaa aaaaatgaat aa 582

<210> 11

<211> 193

<212> PRT

<213> Crenarchaeote

<400> 11

Met Gln Leu Glu Asn Asn Asn Ile Gly Glu Glu Lys Asn Ser Lys Asn
1 5 10 15

Thr Leu Ser Glu Glu Ala Gly Leu Gln Ser Val Phe Glu Asn Phe Ile
20 25 30

Lys Gln Leu Thr Glu Leu Asn Ser Leu Thr Thr Leu Gly Pro Phe Thr
35 40 45

Ser Leu Met Asn Asp Pro Asn Leu Asn Leu Asn Thr Leu Lys Glu His
50 55 60

Gly Asn Leu Leu Leu Arg Tyr Gln Ser Phe Leu Asn Leu Tyr Phe Ser
65 70 75 80

Arg Met Ile Asn Ala Tyr Leu Leu Ala Val Asn Lys Val Ser Ser Ala
85 90 95

Ile Asp Glu Lys Asn Pro Asp Asp Ile Arg Lys Ile Ile Ile Asn Thr
100 105 110

Phe Glu Asp Val Phe Ser Ser Met Leu Gln Ser Thr Asp Phe Ser Ile
115 120 125

Asn Tyr Asn Asn Leu Leu Asn Ser Ser Ile Asp Val Ile Lys Ser Tyr
130 135 140

Gln Lys Ile Tyr Asp Ser Asn Ala Val Leu Phe Arg Ser Gln Gln Gln
145 150 155 160

Leu Ser Lys Glu Glu Lys Asp Leu Leu Phe Tyr Asn Leu Tyr Glu Ile
165 170 175

Lys Lys Ile Ser Leu Glu Ile Lys Lys Lys Leu Asn Glu Lys Lys Asn
180 185 190

Glu

<210> 12

<211> 438

<212> DNA

<213> Crenarchaeote

<400> 12

```

atgcctacaa gttcagatgt ttatatacatg tccaaaccag cggtggtatg tatacattct      60
tttgacatgg tgggcgggta tgcgcataacc caaaaactaa gatgctgtat cagcctcggg      120
aagagggtta tgtgggggac aatagaaaga atccatccac aaacgaatgg ttttggcaaa      180
tgtctgcctt ggctcattgt ttctatatat ggtttcgcca tagataatat ttgggtaatt      240
actcatgtgc ctgtatacaa aaatagacaa ccatctctac ctatatataa attttttgac      300
aagggtcagg ttctttttct ctttttttg caaattatgc cgggccaccc agaacaacaa      360
cccgctttg ggcccgcaat gattactggg caaccctaat ctagcgcccc acgccagga      420
gtgcaagtga caatgtga                                     438

```

<210> 13

<211> 145

<212> PRT

<213> Crenarchaeote

<400> 13

```

Met Pro Thr Ser Ser Asp Val Leu Tyr Met Ser Lys Pro Ala Val Val
1           5           10          15

Cys Ile His Ser Phe Asp Met Val Gly Gly Tyr Ala His Thr Gln Lys
          20           25           30

Leu Arg Cys Cys Ile Ser Leu Gly Lys Arg Val Met Trp Gly Thr Ile
          35           40           45

Glu Arg Ile His Pro Gln Thr Asn Gly Phe Gly Lys Cys Leu Pro Trp
          50           55           60

Leu Ile Val Ser Ile Tyr Gly Phe Ala Ile Asp Asn Ile Trp Val Ile
65           70           75           80

```

Thr His Val Pro Val Tyr Lys Asn Arg Gln Pro Ser Leu Pro Ile Tyr
85 90 95

Lys Phe Phe Asp Lys Gly Gln Val Leu Phe Leu Leu Phe Leu Gln Ile
100 105 110

Met Pro Gly His Pro Glu Thr Asn Pro Ala Phe Gly Pro Ala Met Ile
115 120 125

Thr Gly Gln Pro Lys Ser Ser Ala Pro Arg Pro Gly Val Gln Val Thr
130 135 140

Met
145

<210> 14

<211> 915

<212> DNA

<213> Crenarchaeote

<400> 14

ttggattctt ggggcgaatc taatattgtc ctttggctgc tacttaggct attcaagcca	60
aaaaccaaga tctttgttgt gtttcacac catgaaccgc ggatatctat ttgcaagaac	120
tattttgaat ttctgtacaa ttatctaata caaaaggcta ctgcggtgat gcttaaggat	180
tctgatatga ttttgaccgt gagtcaagcg tcaaagcatg aactcaacac agtctatgga	240
ataggggtta gcaaaatcaa taatttgaag gaaacagcaa ataaaaaac cagggaatta	300
gcaaaaaatc tgaccaacag aattgccatt gtaggaactg gaatagataa aaatatcttt	360
ttaaaggatt ccaacagagg agtaatcaac aataaaaagg acattgattt tctttgtatc	420
ggaaggatag aaaaatttca tggactggag gaaatttga ctgcaataaa aacactcaga	480
ccagaatcta attttgtaat ggttgggcgc ataccctcg ataaggctgc aaaactacgt	540
aatgcgggta tagatcacag aggctttgtc tccgaggaag aaaagattag cctttattct	600
aaatctaaag tctttatttt tccatcatcc agagagggtt ttggcattgc tgtggctgag	660
gccttagttt cgtgtgttcc cactgttgcc tggaaactcc ccgtttttga agaactatac	720
ttaaaaaatg gtaatacaaa cataaaacta atagaatatg gagaaaccac cctgtttgca	780
gaagagtgcg taaaaatgct aaataaatat ggcataatca aaaaggcgac tgaaggaaaa	840
aaggtcagtt tccaactccc aaactggcag acagtggcaa aaaatgtaat gacaacaata	900

gaatctgtaa cctaa

915

<210> 15

<211> 304

<212> PRT

<213> Crenarchaeote

<400> 15

Met Asp Ser Trp Gly Glu Ser Asn Ile Val Leu Trp Leu Leu Arg
1 5 10 15

Leu Phe Lys Pro Lys Thr Lys Ile Phe Val Val Phe His His His Glu
20 25 30

Pro Arg Ile Ser Ile Cys Lys Asn Tyr Phe Glu Phe Leu Tyr Asn Tyr
35 40 45

Leu Ile Gln Lys Ala Thr Ala Val Met Leu Lys Asp Ser Asp Met Ile
50 55 60

Leu Thr Val Ser Gln Ala Ser Lys His Glu Leu Asn Thr Val Tyr Gly
65 70 75 80

Ile Gly Val Ser Lys Ile Asn Asn Leu Lys Glu Thr Ala Asn Lys Lys
85 90 95

Thr Arg Glu Leu Ala Lys Asn Leu Thr Asn Arg Ile Ala Ile Val Gly
100 105 110

Thr Gly Ile Asp Lys Asn Ile Phe Leu Lys Asp Ser Asn Arg Gly Val
115 120 125

Ile Asn Asn Lys Lys Asp Ile Asp Phe Leu Cys Ile Gly Arg Ile Glu
130 135 140

Lys Phe His Gly Leu Glu Glu Ile Trp Thr Ala Ile Lys Thr Leu Arg
145 150 155 160

Pro Glu Ser Asn Phe Val Met Val Gly Arg Ile Pro Pro Asp Lys Ala
165 170 175

Ala Lys Leu Arg Asn Ala Gly Ile Asp His Arg Gly Phe Val Ser Glu

180

185

190

Glu Glu Lys Ile Ser Leu Tyr Ser Lys Ser Lys Val Phe Ile Phe Pro

195

200

205

Ser Ser Arg Glu Gly Phe Gly Ile Ala Val Ala Glu Ala Leu Val Ser

210

215

220

Cys Val Pro Thr Val Ala Trp Lys Leu Pro Val Phe Glu Glu Leu Tyr

225

230

235

240

Leu Lys Asn Gly Asn Thr Asn Ile Lys Leu Ile Glu Tyr Gly Glu Thr

245

250

255

Thr Leu Phe Ala Glu Glu Cys Val Lys Met Leu Asn Lys Tyr Gly Ile

260

265

270

Ile Lys Lys Ala Thr Glu Gly Lys Lys Val Ser Phe Gln Leu Pro Asn

275

280

285

Trp Gln Thr Val Ala Lys Asn Val Met Thr Thr Ile Glu Ser Val Thr

290

295

300

<210> 16

<211> 1692

<212> DNA

<213> Crenarchaeote

<400> 16

atgtgtggaa ttgttggaat tttaagtaaa aaagagagaa atgttgcccc cttgatagga 60

aaaatgctat cctgtatgaa aaaccggggt ccgatggca tgggtttgtc tacagagaat 120

caaatagttt attctgatac ctttgataat ccattgtttt cacaggtaga ggggcatgac 180

gttttaggtc acagtcgttt ggcaatagtt ggtggctcct gtggtcagca gccgtttgtg 240

agttgtgata aaaaactcat tctggagcat aatggtgaaa tatataacta taaagaaatc 300

agaaagaacc tttctgcaca tcacactttt actacctcga ctgatagtga agttattgtt 360

caccttcttg aagaccatta tcaaaacact aaaggcgatc taatcgaagc tataaggaga 420

accgttacct agcttgatgg aatttatgtt ttggcgatta gagagcagtc cacaggagat 480

attgtgctgg tacgggatgg cattggagta agacaaattt actatggtga aagtagtgat 540
 ttcattgcat ttgcatcaga aagaaaagcc ttatggaaaa ttgctatgtc cgaccaaadc 600
 aaaagacttt tgccaggcta tgctcttgtc atttcgcgga aggaagggtc ctccaatttc 660
 aagactacat tgtttccgat ttctgtaaata acaaaaaaat caatatgtga gaaatattca 720
 atcctgtaca cagacatcga ttctgcggtt aacgcatatg gtgatacatt ggttgaatct 780
 atgagaaaac gtgtgagtga ctttaaaaaa atcgggtattg ttttctccgg tgggattgac 840
 agtgtaattg tagcgtattt ggcaaaacaa atggcccccg aagttatttg ctatacgtct 900
 gggattaaag gttcaagtga taccctcaac tcacttgaga tagcagaaaa acttgacctc 960
 aagttggaaa tagaacagat gactgaaagt gatgttgaaa gtaccattcc aaaaataatc 1020
 agcataattg aagatgacaa catgggacag gttgaggttg ccattccaat atatggcgcg 1080
 gttaaattgg ctacgaaca gggaatacgg gtaatgctta caggtcaggg ggcagacgaa 1140
 ctgtttgccg gatattcctg gtattccaaa attgttaaaa aacacggata cgaaaaaatt 1200
 cagggtatcc tgatagagga cattaagtta ctttacaag aacactgga aagagaggac 1260
 aaaataacca tgtctcaaag catagagtta cgcgaaccct ttttagatac taatctgata 1320
 gacacggtac tgagaataga tccgcgactc aatattcaaa acaatggcaa taactatgac 1380
 aacctaggaa aaagggttca ccgcaaactt gcagaaaaac tagggattcc aaaagagata 1440
 gcgtatagaa taaaggaagc agctcagcat gggtctggga tacacaacac cctcaatact 1500
 ttggccatga aaaatggttt tacggaatcc aagggttaatt ctagtattct ggacaaattg 1560
 aaaaaagggt agcttatcgg cagctcacia agatatgggc atctttttga aaaggaacaa 1620
 atctggagtt tggagccgca tatacagatg tatttggaga atatttcaaa aaacatattg 1680
 ccaaggaact ga 1692

<210> 17

<211> 563

<212> PRT

<213> Crenarchaeote

<400> 17

Met Cys Gly Ile Val Gly Ile Leu Ser Lys Lys Glu Arg Asn Val Ala
 1 5 10 15

Pro Leu Ile Gly Lys Met Leu Ser Cys Met Lys Asn Arg Gly Pro Asp
 20 25 30

Gly Met Gly Leu Ser Thr Glu Asn Gln Ile Val Tyr Ser Asp Thr Phe
 35 40 45

Asp Asn Pro Leu Phe Ser Gln Val Glu Gly His Asp Val Leu Gly His
 50 55 60

Ser Arg Leu Ala Ile Val Gly Gly Ser Cys Gly Gln Gln Pro Phe Val
 65 70 75 80

Ser Cys Asp Lys Lys Leu Ile Leu Glu His Asn Gly Glu Ile Tyr Asn
 85 90 95

Tyr Lys Glu Ile Arg Lys Asn Leu Ser Ala His His Thr Phe Thr Thr
 100 105 110

Ser Thr Asp Ser Glu Val Ile Val His Leu Leu Glu Asp His Tyr Gln
 115 120 125

Asn Thr Lys Gly Asp Leu Ile Glu Ala Ile Arg Arg Thr Val Thr Gln
 130 135 140

Leu Asp Gly Ile Tyr Val Leu Ala Ile Arg Glu Gln Ser Thr Gly Asp
 145 150 155 160

Ile Val Leu Val Arg Asp Gly Ile Gly Val Arg Gln Ile Tyr Tyr Gly
 165 170 175

Glu Ser Ser Asp Phe Ile Ala Phe Ala Ser Glu Arg Lys Ala Leu Trp
 180 185 190

Lys Ile Ala Met Ser Asp Gln Ile Lys Arg Leu Leu Pro Gly Tyr Ala
 195 200 205

Leu Val Ile Ser Arg Lys Glu Gly Ser Ser Asn Phe Lys Thr Thr Leu
 210 215 220

Phe Pro Ile Ser Val Asn Thr Lys Lys Ser Ile Cys Glu Lys Tyr Ser
 225 230 235 240

Ile Leu Tyr Thr Asp Ile Asp Ser Ala Val Asn Ala Tyr Gly Asp Thr
 245 250 255

Leu Val Glu Ser Met Arg Lys Arg Val Ser Asp Phe Lys Lys Ile Gly
260 265 270

Ile Val Phe Ser Gly Gly Ile Asp Ser Val Ile Val Ala Tyr Leu Ala
275 280 285

Lys Gln Met Ala Pro Glu Val Ile Cys Tyr Thr Ser Gly Ile Lys Gly
290 295 300

Ser Ser Asp Ile Leu Asn Ser Leu Glu Ile Ala Glu Lys Leu Asp Leu
305 310 315 320

Lys Leu Glu Ile Glu Gln Met Thr Glu Ser Asp Val Glu Ser Thr Ile
325 330 335

Pro Lys Ile Ile Ser Ile Ile Glu Asp Asp Asn Met Gly Gln Val Glu
340 345 350

Val Ala Ile Pro Ile Tyr Gly Ala Val Lys Leu Ala His Glu Gln Gly
355 360 365

Ile Arg Val Met Leu Thr Gly Gln Gly Ala Asp Glu Leu Phe Gly Gly
370 375 380

Tyr Ser Trp Tyr Ser Lys Ile Val Lys Lys His Gly Tyr Glu Lys Ile
385 390 395 400

Gln Gly Tyr Leu Ile Glu Asp Ile Lys Leu Leu Tyr Lys Glu Thr Leu
405 410 415

Glu Arg Glu Asp Lys Ile Thr Met Ser Gln Ser Ile Glu Leu Arg Glu
420 425 430

Pro Phe Leu Asp Thr Asn Leu Ile Asp Thr Val Leu Arg Ile Asp Pro
435 440 445

Arg Leu Asn Ile Gln Asn Asn Gly Asn Asn Tyr Asp Asn Leu Gly Lys
450 455 460

Arg Val His Arg Lys Leu Ala Glu Lys Leu Gly Ile Pro Lys Glu Ile
465 470 475 480

Ala Tyr Arg Ile Lys Glu Ala Ala Gln His Gly Ser Gly Ile His Asn
485 490 495

Thr Leu Asn Thr Leu Ala Met Lys Asn Gly Phe Thr Glu Ser Lys Val
500 505 510

Asn Ser Ser Tyr Leu Asp Lys Leu Lys Lys Arg Glu Leu Ile Gly Ser
515 520 525

Ser Gln Arg Tyr Gly His Leu Phe Glu Lys Glu Gln Ile Trp Ser Leu
530 535 540

Glu Pro His Ile Gln Met Tyr Leu Glu Asn Ile Ser Lys Asn Ile Leu
545 550 555 560

Pro Arg Asn

<210> 18

<211> 666

<212> DNA

<213> Crenarchaeote

<400> 18
ttgttgatc ctatggaatt taaatctaca ttggccgttt ttgatatgga tgggacgcta 60
attgatggaa ggctaattga ggtattgtca aaaaagtttg gcttgatgc tcaggtcaga 120
cacatccagt ccgacaaatc cattccaggc tatgttaaga cacagaagat agccgctgtg 180
attaggggaa tagaagaaag ggaaatagaa attgctttgg actccatccc cctgcaaag 240
aacagccagg aggtgatatc tttgctgaag aaaaaagggc tcagaatagg gataattaca 300
gatagttaca gtgttgctgc tcaggccttg gtgaacaaac ttgatttgga ctttttttat 360
gcaaataaat tgaaggtaga caatgggata gtcaccggag aaataaatat gccgttagga 420
tgggaaaaaa tagactgttt ttgcaagaat tctgtgtgta agagatatca catggaaatc 480
catgcaaaga aaatctgtgc agacataaaa aatacaattg ctattggcga tactaaaggt 540
gacctgtgca tgataaagca ggcaggaata ggtatcgcat atatgcctaa ggataaatat 600
ataaatgaaa caataaataa ggtaaacaca ccggatatga ttggtgtcct tgattttata 660
gagtag 666

<210> 19

<211> 221

<212> PRT

<213> Crenarchaeote

<400> 19

Met Leu Tyr Pro Met Glu Phe Lys Ser Thr Leu Ala Val Phe Asp Met
1 5 10 15

Asp Gly Thr Leu Ile Asp Gly Arg Leu Ile Glu Val Leu Ser Lys Lys
20 25 30

Phe Gly Leu Tyr Ala Gln Val Arg His Ile Gln Ser Asp Lys Ser Ile
35 40 45

Pro Gly Tyr Val Lys Thr Gln Lys Ile Ala Ala Val Ile Arg Gly Ile
50 55 60

Glu Glu Arg Glu Ile Glu Ile Ala Leu Asp Ser Ile Pro Pro Ala Lys
65 70 75 80

Asn Ser Gln Glu Val Ile Ser Leu Leu Lys Lys Lys Gly Phe Arg Ile
85 90 95

Gly Ile Ile Thr Asp Ser Tyr Ser Val Ala Ala Gln Ala Leu Val Asn
100 105 110

Lys Leu Asp Leu Asp Phe Phe Tyr Ala Asn Glu Leu Lys Val Asp Asn
115 120 125

Gly Ile Val Thr Gly Glu Ile Asn Met Pro Leu Gly Trp Glu Lys Ile
130 135 140

Asp Cys Phe Cys Lys Asn Ser Val Cys Lys Arg Tyr His Met Glu Ile
145 150 155 160

His Ala Lys Lys Ile Cys Ala Asp Ile Lys Asn Thr Ile Ala Ile Gly
165 170 175

Asp Thr Lys Gly Asp Leu Cys Met Ile Lys Gln Ala Gly Ile Gly Ile
180 185 190

Ala Tyr Met Pro Lys Asp Lys Tyr Ile Asn Glu Thr Ile Asn Lys Val
195 200 205

Asn Thr Pro Asp Met Ile Gly Val Leu Asp Phe Ile Glu
 210 215 220

<210> 20

<211> 1212

<212> DNA

<213> Crenarchaeote

<400> 20

```

atgagattag attatccacc taactatacc gagaggatag gagcagttag tatccatgcg      60
cttcaaaaga tttatgagat cgattccgga aagatgccca agtttaatgg cctgcatcag      120
catcagtcta taaaggcctt tggttatgac gaactgtcaa gcatattcca agaacttgcc      180
atagtcattc cagtaaagaa cgaaaaaatc agccttcttg aaggagtatt gagcggattt      240
ccaaatgaat gtctcatcat catagtttcc aatagccaaa ggactcctgt cgacagattt      300
gccatggagg ttgaaatggg aaggcagtac tctagttttg cagacaagaa aataatgatt      360
attcaccaaa atgatcctga gctggctaata acttttaaga aaataaagta tagatccatc      420
ctcaacacca aaagtcaggt tcgtagtgga aaggctgaag gaatgataat tggaatattg      480
ctggcaaaaa tgcacctaaa agagtacatt ggatttattg acagtgataa ttattttcca      540
ggagcagtaa atgaatatgt caagatcttt gcagcgggat ttggaatggc aaccacccca      600
tacagcaata tcagaatata gtggcggttc aaacccaaaa tcgtaaacaa ctactacaa      660
ttccaagat ggggtagaat ttcagaatcc agtaacaaat acctgaacgc tctaatatcc      720
cacatcacag ggtttgaaag ggagattatc acgactggaa atgcagggtga gcatgcatta      780
tccatgtccc ttgcagaaaa tctcaactat tcaagcggat attcggttga gccctatgag      840
tttatcaaca ttttagaaaa gtttggagggt ctactcccat caaacaatcc tgacatcata      900
gaaaagggtg tcgaaatatt tcaaatagag accaggaatc cacactttca tgaggaaaaa      960
ggaaatgatc atttggcagg catgatgcaa gaatctcttc tcgcaataaa caacagcaaa     1020
atttgaaca cagaactgac cagggaataa aatgaccatt tactcatgct tcaggtaaaa     1080
cacaataatg atatgaccaa actcaacttt aagaaaaaac accttataat ggatcccata     1140
aaaataatac ccatcgacaa attcgccgaa tttgtagtta agaattctaa aaccttcatt     1200
agaattggat aa                                                    1212

```

<210> 21

<211> 403

<212> PRT

<213> Crenarchaeote

<400> 21

Met Arg Leu Asp Tyr Pro Pro Asn Tyr Thr Glu Arg Ile Gly Ala Val
1 5 10 15

Ser Ile His Ala Leu Gln Lys Ile Tyr Glu Ile Asp Ser Gly Lys Met
20 25 30

Pro Lys Phe Asn Gly Leu His Gln His Gln Ser Ile Lys Ala Phe Gly
35 40 45

Tyr Asp Glu Leu Ser Ser Ile Phe Gln Glu Leu Ala Ile Val Ile Pro
50 55 60

Val Lys Asn Glu Lys Ile Ser Leu Leu Glu Gly Val Leu Ser Gly Ile
65 70 75 80

Pro Asn Glu Cys Leu Ile Ile Ile Val Ser Asn Ser Gln Arg Thr Pro
85 90 95

Val Asp Arg Phe Ala Met Glu Val Glu Met Val Arg Gln Tyr Ser Ser
100 105 110

Phe Ala Asp Lys Lys Ile Met Ile Ile His Gln Asn Asp Pro Glu Leu
115 120 125

Ala Asn Thr Phe Lys Lys Ile Lys Tyr Arg Ser Ile Leu Asn Thr Lys
130 135 140

Ser Gln Val Arg Ser Gly Lys Ala Glu Gly Met Ile Ile Gly Ile Leu
145 150 155 160

Leu Ala Lys Met His Leu Lys Glu Tyr Ile Gly Phe Ile Asp Ser Asp
165 170 175

Asn Tyr Phe Pro Gly Ala Val Asn Glu Tyr Val Lys Ile Phe Ala Ala
180 185 190

Gly Phe Gly Met Ala Thr Thr Pro Tyr Ser Asn Ile Arg Ile Ser Trp
195 200 205

Arg Ser Lys Pro Lys Ile Val Asn Asn Ser Leu Gln Phe Pro Arg Trp
210 215 220

Gly Arg Ile Ser Glu Ser Ser Asn Lys Tyr Leu Asn Ala Leu Ile Ser
225 230 235 240

His Ile Thr Gly Phe Glu Arg Glu Ile Ile Thr Thr Gly Asn Ala Gly
245 250 255

Glu His Ala Leu Ser Met Ser Leu Ala Glu Asn Leu Asn Tyr Ser Ser
260 265 270

Gly Tyr Ser Val Glu Pro Tyr Glu Phe Ile Asn Ile Leu Glu Lys Phe
275 280 285

Gly Gly Leu Leu Pro Ser Asn Asn Pro Asp Ile Ile Glu Lys Gly Ile
290 295 300

Glu Ile Phe Gln Ile Glu Thr Arg Asn Pro His Phe His Glu Glu Lys
305 310 315 320

Gly Asn Asp His Leu Ala Gly Met Met Gln Glu Ser Leu Leu Ala Ile
325 330 335

Asn Asn Ser Lys Ile Cys Asn Thr Glu Leu Thr Arg Glu Ile Asn Asp
340 345 350

His Leu Leu Met Leu Gln Val Lys His Asn Asn Asp Met Thr Lys Leu
355 360 365

Asn Phe Lys Lys Lys His Leu Ile Met Asp Pro Ile Lys Ile Ile Pro
370 375 380

Ile Asp Lys Phe Ala Glu Phe Val Val Lys Asn Ser Lys Thr Phe Ile
385 390 395 400

Arg Ile Gly

<210> 22

<211> 1164

<212> DNA

<213> Crenarchaeote

<400> 22
 atgagtgatg ctatcgaaaa tgtcctgatc cttcagggag gaggatcttt ggggtgcattt 60
 gggttgcgggg tctacaaagc actagtaaac aataacataa aacttgatat cctgtctggc 120
 acatcaattg gcgggttgaa tgccacagtt attgccggca gtaaagaaga tcgtccagaa 180
 aaatcattgg agaatttttg gatggaaata gctgatacta ataatggtaa tattaatata 240
 taccttaatt tccccctttt tgaaagtccc tttcctgggc aaattccttt ccccttggca 300
 tcagaatcaa cactatcatt ctacagctct gccatttatg gaaatagaaa aatctttctg 360
 ccaagatggg gacctgaaaa tatctttaaa gatccacagt atttcacacc tagcaaatgg 420
 acatatttgt atgaccattc acctttggta aaaaccttgg aaaagtacat tgattatagc 480
 aaattacagc caaacggtaa gccaacgca aggctaataa taaccgcagt taacgtgatg 540
 acggcggagc cccttatctt tgacagtgcc aagcaacaaa taaccccaaa acacatactt 600
 gcaaccactg cctatccaac atattttttt caatgggtgg aattggaaaa agggcttttt 660
 gcctgggatg gaagtttact aagcaatacc ccgctaagag aagtaataga cgcacgccc 720
 gcaaaggaca aaagaatctt tcttgtcgag aactatccta aaaatattga aaagcttccg 780
 tcaaacctac aggaagtcaa gcatagggca agagacataa tgttcagcga caagaccgtc 840
 cacagtatac acatgtocaa agcaattacc cttcaactta agcttattga tgatctgtat 900
 aaaatgctag agtattactt taattcagaa aaaatcgagg aaaaggagaa gtttgaaaaa 960
 attcgtgcga gatacaaaaa agtttcagaa gaacacggcg cagagattaa aggtgtctac 1020
 tatataacac gggacgagcc atccccctcc ctttatgaga atgcagactt ttcaaaaaat 1080
 gcaataaagg catcgattaa tgatggagaa caaaaggctg acaggataat aaaagaaatc 1140
 caaacgaaag gaaaacgaaa ataa. 1164

<210> 23

<211> 387

<212> PRT

<213> Crenarchaeote

<400> 23

Met Ser Asp Ala Ile Glu Asn Val Leu Ile Leu Gln Gly Gly Gly Ser
 1 5 10 15

Leu Gly Ala Phe Gly Cys Gly Val Tyr Lys Ala Leu Val Asn Asn Asn
20 25 30

Ile Lys Leu Asp Ile Leu Ser Gly Thr Ser Ile Gly Gly Leu Asn Ala
35 40 45

Thr Val Ile Ala Gly Ser Lys Glu Asp Arg Pro Glu Lys Ser Leu Glu
50 55 60

Asn Phe Trp Met Glu Ile Ala Asp Thr Asn Asn Gly Asn Ile Asn Thr
65 70 75 80

Tyr Leu Asn Phe Pro Phe Phe Glu Ser Pro Phe Pro Gly Gln Ile Pro
85 90 95

Phe Pro Leu Ala Ser Glu Ser Thr Leu Ser Phe Tyr Ser Ser Ala Ile
100 105 110

Tyr Gly Asn Arg Lys Ile Phe Leu Pro Arg Trp Gly Pro Glu Asn Ile
115 120 125

Phe Lys Asp Pro Gln Tyr Phe Thr Pro Ser Lys Trp Thr Tyr Leu Tyr
130 135 140

Asp His Ser Pro Leu Val Lys Thr Leu Glu Lys Tyr Ile Asp Tyr Ser
145 150 155 160

Lys Leu Gln Pro Asn Gly Lys Pro Asn Ala Arg Leu Ile Ile Thr Ala
165 170 175

Val Asn Val Met Thr Ala Glu Pro Leu Ile Phe Asp Ser Ala Lys Gln
180 185 190

Gln Ile Thr Pro Lys His Ile Leu Ala Thr Thr Ala Tyr Pro Thr Tyr
195 200 205

Phe Phe Gln Trp Val Glu Leu Glu Lys Gly Leu Phe Ala Trp Asp Gly
210 215 220

Ser Leu Leu Ser Asn Thr Pro Leu Arg Glu Val Ile Asp Ala Ser Pro
225 230 235 240

Ala Lys Asp Lys Arg Ile Phe Leu Val Glu Asn Tyr Pro Lys Asn Ile
245 250 255

Glu Lys Leu Pro Ser Asn Leu Gln Glu Val Lys His Arg Ala Arg Asp
260 265 270

Ile Met Phe Ser Asp Lys Thr Val His Ser Ile His Met Ser Lys Ala
275 280 285

Ile Thr Leu Gln Leu Lys Leu Ile Asp Asp Leu Tyr Lys Met Leu Glu
290 295 300

Tyr Tyr Phe Asn Ser Glu Lys Ile Glu Glu Lys Glu Lys Phe Glu Lys
305 310 315 320

Ile Arg Ala Arg Tyr Lys Lys Val Ser Glu Glu His Gly Ala Glu Ile
325 330 335

Lys Gly Val Tyr Tyr Ile Thr Arg Asp Glu Pro Ser Pro Ser Leu Tyr
340 345 350

Glu Asn Ala Asp Phe Ser Lys Asn Ala Ile Lys Ala Ser Ile Asn Asp
355 360 365

Gly Glu Gln Lys Ala Asp Arg Ile Ile Lys Glu Ile Gln Thr Lys Gly
370 375 380

Lys Arg Lys
385

<210> 24

<211> 882

<212> DNA

<213> Crenarchaeote

<400> 24

atggaactta acgcagcagt aattgtgaaa ctcgagccgg atttttctga agggaatgta 60

agctataatt cagacggaac acttaacaga gcagaaacaa aaaacatttt ggggccccat 120

agcgcagcag catccctagc agccctgtac tcaaaagtaa aacatggaac gcatgtttct 180

gtgggcacaa tgggtcctcc aatagcagaa tcggccttac agcaatctca actgatttgc 240

gacgctgatg aactgcatct ttatagtgat cgcacttttg caggagccga caccctggcc 300

acagctgaag ttttgatagc aggaataaaa aaaatggcaa atggtcaaga tgtggacatt 360

gttttctcag ggcacagggc atctgatggc gaaacagggc aaacaggacc ccagacagca 420
tggaattag gttatccgtt ccttggaat gttattgatt acgatattga cgttggtgaag 480
agaattgtaa ggttacaacg tctaatacag atttacggtc atcctgatat tatagaggag 540
atggaggcgc ctctaccggt ttttatcaca ctggacccat cctacaatcc gtcttttaac 600
acggtatccc aaaggctcag actagcacga aacctacagg aagcccatga tagatcacia 660
aggataagg aatatctcaa aactttcaat gccatggaac tagaagtcaa tccaaagtct 720
gtcggactgc ctggctctcc caccatagtt tataaagttg aaaaaatacc aagggcaaag 780
gcaaatagaa aagcagatgt tgtggatggg tctaaccagg atagtctaag gcaggttgca 840
cgccgaatcc atgatgtttt agggggtgta gtcataaagt ga 882

<210> 25

<211> 293

<212> PRT

<213> Crenarchaeote

<400> 25

Met Glu Leu Asn Ala Ala Val Ile Val Lys Leu Glu Pro Asp Phe Ser
1 5 10 15

Glu Gly Asn Val Ser Tyr Asn Ser Asp Gly Thr Leu Asn Arg Ala Glu
20 25 30

Thr Lys Asn Ile Leu Gly Pro His Ser Ala Ala Ala Ser Leu Ala Ala
35 40 45

Leu Tyr Ser Lys Val Lys His Gly Thr His Val Ser Val Gly Thr Met
50 55 60

Gly Pro Pro Ile Ala Glu Ser Ala Leu Gln Gln Ser Gln Leu Ile Cys
65 70 75 80

Asp Ala Asp Glu Leu His Leu Tyr Ser Asp Arg Ile Phe Ala Gly Ala
85 90 95

Asp Thr Leu Ala Thr Ala Glu Val Leu Ile Ala Gly Ile Lys Lys Met
100 105 110

Ala Asn Gly Gln Asp Val Asp Ile Val Phe Ser Gly His Arg Ala Ser
115 120 125

Asp Gly Glu Thr Gly Gln Thr Gly Pro Gln Thr Ala Trp Lys Leu Gly
130 135 140

Tyr Pro Phe Leu Gly Asn Val Ile Asp Tyr Asp Ile Asp Val Val Lys
145 150 155 160

Arg Ile Val Arg Val Gln Arg Leu Ile Lys Ile Tyr Gly His Pro Asp
165 170 175

Ile Ile Glu Glu Met Glu Ala Pro Leu Pro Val Phe Ile Thr Leu Asp
180 185 190

Pro Ser Tyr Asn Pro Ser Phe Asn Thr Val Ser Gln Arg Leu Arg Leu
195 200 205

Ala Arg Asn Leu Gln Glu Ala His Asp Arg Ser Gln Arg Tyr Lys Glu
210 215 220

Tyr Leu Lys Thr Phe Asn Ala Met Glu Leu Glu Val Asn Pro Lys Ser
225 230 235 240

Val Gly Leu Pro Gly Ser Pro Thr Ile Val Tyr Lys Val Glu Lys Ile
245 250 255

Pro Arg Ala Lys Ala Asn Arg Lys Ala Asp Val Val Asp Gly Ser Asn
260 265 270

Gln Asp Ser Leu Arg Gln Val Ala Arg Arg Ile His Asp Val Leu Gly
275 280 285

Gly Val Val Ile Lys
290

<210> 26

<211> 1284

<212> DNA

<213> Crenarchaeote

<400> 26

gtgacatcat cactatctgc catacctgac gctaaactag acgaaaggcc aaacccaaat

60

gcccattgta atgacaaccc agaaaaagaa aggggagaca acaacaggca tctgtatgtt 120
gtgatagaac aagaggaagg caccatatta cctgtgagtt ttgaaatgct tggtaggca 180
agaaggctaa tggatgattt taatcacaaa tacaagccag aggaaaaagt ggttgcgatt 240
atactcggcc ataacatcaa gcacctgtgc caggaactaa tccaccatgg tgcagacgca 300
gtgatttatg ccgaccaccc ggagctccgc caccaagaa atcttcttta tacaaaggtt 360
gtctgccaaa ttgctacgga caaagagagc gccgccagaa tttggccatc aaatcccgat 420
tttaacagac cccgttacat gtttttttcc gcagatgaca caggaaggca tttatcatca 480
accgttttgg cagaattgca atcagggtcg gcacagaca taaacaaact tgttatcaat 540
gatttagaaa taaggcatga acacaagaca aagggtaaac ccattgtcta tgaaaagaca 600
cttgaaatgt acagaccaga cttttcaggc tttctttgga ccaccatact ctgcttggat 660
aatataaatc ccgagaacag aaggaaattc catccacagg catgcagtat aatcccaggc 720
gtctttcccc aaatggaagg agatacggat agaaagggtta ccataataga gttcagccca 780
accatagccc aggaagacct tagaataaaa ataatcaaca gaagagtaat caaaagcaaa 840
gtcgatttta gcaataaaaa aataatcggt agttttggaa ggggaataaa ggagtctccc 900
gaacaaaaca taaaactgat agagaacctt gcaaaggaaa tagaagcaga aataggaata 960
tactgccc tttcaaagaa accctatcca ataagcgaaa gtctgtcgtc aacctatatg 1020
attcctgaca gggttatcgg cacaagcgga agaaaggtaa atcctcaggt gtattttgca 1080
ataggaataa gcggggctgt ccaacacata gcggggatga aagaatcgga atttgtgatt 1140
tccatcaatc cagacagtga agctcccata atagatgaat ccgatgtttt aatcaaagga 1200
aaaatcgagc aggtgctgcc tctcctgata aatgaattaa aaaaatacaa agagagactg 1260
caaataccac aggagataga atga 1284

<210> 27

<211> 427

<212> PRT

<213> Crenarchaeote

<400> 27

Met Thr Ser Ser Leu Ser Ala Ile Pro Asp Ala Lys Leu Asp Glu Arg
1 5 10 15

Pro Asn Gln Asn Ala His Val Asn Asp Asn Pro Glu Lys Glu Arg Gly
20 25 30

Asp Asn Asn Arg His Leu Tyr Val Val Ile Glu Gln Glu Glu Gly Thr
35 40 45

Ile Leu Pro Val Ser Phe Glu Met Leu Gly Glu Ala Arg Arg Leu Met
50 55 60

Asp Asp Phe Asn His Lys Tyr Lys Pro Glu Glu Lys Val Val Ala Ile
65 70 75 80

Ile Leu Gly His Asn Ile Lys His Leu Cys Gln Glu Leu Ile His His
85 90 95

Gly Ala Asp Ala Val Ile Tyr Ala Asp His Pro Glu Leu Arg His Pro
100 105 110

Arg Asn Leu Leu Tyr Thr Lys Val Val Cys Gln Ile Ala Thr Asp Lys
115 120 125

Glu Ser Ala Ala Arg Ile Trp Pro Ser Asn Pro Asp Phe Asn Arg Pro
130 135 140

Arg Tyr Met Phe Phe Ser Ala Asp Asp Thr Gly Arg His Leu Ser Ser
145 150 155 160

Thr Val Leu Ala Glu Leu Gln Ser Gly Leu Ala Ser Asp Ile Asn Lys
165 170 175

Leu Val Ile Asn Asp Leu Glu Ile Arg His Glu His Lys Thr Lys Gly
180 185 190

Lys Pro Ile Val Tyr Glu Lys Thr Leu Glu Met Tyr Arg Pro Asp Phe
195 200 205

Ser Gly Phe Leu Trp Thr Thr Ile Leu Cys Leu Asp Asn Ile Asn Pro
210 215 220

Glu Asn Arg Arg Lys Phe His Pro Gln Ala Cys Ser Ile Ile Pro Gly
225 230 235 240

Val Phe Pro Gln Met Glu Gly Asp Thr Asp Arg Lys Gly Thr Ile Ile
245 250 255

Glu Phe Ser Pro Thr Ile Ala Gln Glu Asp Leu Arg Ile Lys Ile Ile
 260 265 270

Asn Arg Arg Val Ile Lys Ser Lys Val Asp Phe Ser Asn Lys Lys Ile
 275 280 285

Ile Val Ser Phe Gly Arg Gly Ile Lys Glu Ser Pro Glu Gln Asn Ile
 290 295 300

Lys Leu Ile Glu Asn Leu Ala Lys Glu Ile Glu Ala Glu Ile Gly Ile
 305 310 315 320

Ser Leu Pro Ile Ser Lys Lys Pro Tyr Pro Ile Ser Glu Ser Leu Ser
 325 330 335

Ser Thr Tyr Met Ile Pro Asp Arg Val Ile Gly Thr Ser Gly Arg Lys
 340 345 350

Val Asn Pro Gln Val Tyr Phe Ala Ile Gly Ile Ser Gly Ala Val Gln
 355 360 365

His Ile Ala Gly Met Lys Glu Ser Glu Phe Val Ile Ser Ile Asn Pro
 370 375 380

Asp Ser Glu Ala Pro Ile Ile Asp Glu Ser Asp Val Leu Ile Lys Gly
 385 390 395 400

Lys Ile Glu Gln Val Leu Pro Leu Leu Ile Asn Glu Leu Lys Lys Tyr
 405 410 415

Lys Glu Arg Leu Gln Ile Pro Gln Glu Ile Glu
 420 425

<210> 28

<211> 1878

<212> DNA

<213> Crenarchaeote

<400> 28

atgacaatgg aaagttttga tgtggcgata attggtggag ggtctgctgg acttgcgcca 60

cttgagcacc tctccaattt gggaaaacag gcaatcctca tagaggcagg aaaaaaata 120

ggaaacaaaa acgtgtctgg gggcatattg tattccaaaa aaacagcaac tggaaaggtc 180

cacaatgtag aagatgtgtt tgataatddd ctggcagacg ctccgctgga aaggaagata 240
ataaaataca tgcttcacgc cgtctcaagg gaaaaagcgt tctctctgga cctgactttg 300
gcacacgact atcaaacgaa ttttgggtac accgtcctgc tcaacaaact actttcatgg 360
tttgcaaggg aagcatctca aagtgcagaa aaactgggtg gagggataat aacaggtgtc 420
catttaaggt cgataatctg gaaagatgac agtaccataa ttatagagac agatgaactt 480
gagccgttcc aggtaaaggc agtcattgca gctgacgggg ttaactcaga ggttgcgcaa 540
ataacaggtg ccagaagcaa gttcacaccg tctgacctct accagggcgt aaaggtgggtg 600
gcaaaattac cagaggggtt gcttgaagag agattcgggg tctcgaaaaa cgagggagcg 660
gctcaccttt tttcaggcga cataacgcta aaccacattg gaggggggtt cctttacaca 720
aacagggaca ccatctcaat tggcgcagta taccattatg actctctaata tgaagagcct 780
acagagccca atgcgctggt caatgcgtta ctgtcaaata cgtttgtgat ggaattgata 840
aaggacgagg ttccaaggat caaggaggac tacagggatc tttcaaagga tgaagaacta 900
aggattaggt tcaaatacaa taaattgata aaaagctgga atgacctaca ccacacatat 960
tattcaccat ctgccgttgc agagcttgtg gcgcagggaa aatacaaata aagggaggag 1020
atcaaggaca aaattgattc attgtacaat gagcttgtaa caaaatacaa cacagaattd 1080
gaaacaaatt acgtggagtt agagtacagc gccaaactgg ttccagatgg aaaaagggtg 1140
agaatgaaaa aaccctactt taaaaacata ttatttgtcg gtgatgctgc gggcaggggc 1200
attdtctctg ggccacgcat agagggcctc aacgtaggca ttgatgacgc ggttagggcc 1260
gcagaagctg tctcaaagtc aatagatcaa aataactttc agtttgacaa cattggtgaa 1320
cgctacacta aatcagtgga tgaaagtcca tataccgcag acatgagcag gatcgacgca 1380
aactatctca aagccgttct tgattgcaca aaaaagggtc ccaaaaacac tcttgggttt 1440
aagtatgggt ctattgtcaa attgatgtca aatagcacct ttaggaatgt atccatagga 1500
attgcaaact ctatagggtg caaaaggctt ttacctgtga ttgagtcaga caaaacctac 1560
aatcaaattc ccatcgagat tgcggagaga aatggcaaag atttgcgga aagctattcc 1620
atagagattc ccaccattgc cgagcgtatt gctaattctga actataatga cgattcactg 1680
tcacacatca aggttttgaa ctcgcaaagt gactttatga aaaaaatgg ccaactgtgc 1740
cctaccaaata gctacagtat tgagaatgag cggataatgc tacagcacga aggatgcata 1800
gagtgtggga catgcgcaag agaaacagaa tggaggcatc ctcgtgggga aaaaggaata 1860
atctataatt acgggtaa 1878

<210> 29

<211> 625

<212> PRT

<213> Crenarchaeote

<400> 29

Met Thr Met Glu Ser Phe Asp Val Ala Ile Ile Gly Gly Gly Ser Ala
1 5 10 15

Gly Leu Ala Ala Leu Glu His Leu Ser Asn Leu Gly Lys Gln Ala Ile
20 25 30

Leu Ile Glu Ala Gly Lys Lys Ile Gly Thr Lys Asn Val Ser Gly Gly
35 40 45

Ile Leu Tyr Ser Lys Lys Thr Ala Thr Gly Lys Val His Asn Val Glu
50 55 60

Asp Val Phe Asp Asn Phe Leu Ala Asp Ala Pro Leu Glu Arg Lys Ile
65 70 75 80

Ile Lys Tyr Met Leu His Ala Val Ser Arg Glu Lys Ala Phe Ser Leu
85 90 95

Asp Leu Thr Leu Ala His Asp Tyr Gln Thr Asn Phe Gly Tyr Thr Val
100 105 110

Leu Leu Asn Lys Leu Leu Ser Trp Phe Ala Arg Glu Ala Ser Gln Ser
115 120 125

Ala Glu Lys Leu Gly Gly Gly Ile Ile Thr Gly Val His Leu Arg Ser
130 135 140

Ile Ile Trp Lys Asp Asp Ser Thr Ile Ile Ile Glu Thr Asp Glu Leu
145 150 155 160

Glu Pro Phe Gln Val Lys Ala Val Ile Ala Ala Asp Gly Val Asn Ser
165 170 175

Glu Val Ala Gln Ile Thr Gly Ala Arg Ser Lys Phe Thr Pro Ser Asp
180 185 190

Leu Tyr Gln Gly Val Lys Val Val Ala Lys Leu Pro Glu Gly Leu Leu
195 200 205

Glu Glu Arg Phe Gly Val Ser Glu Asn Glu Gly Ala Ala His Leu Phe
210 215 220

Ser Gly Asp Ile Thr Leu Asn His Ile Gly Gly Gly Phe Leu Tyr Thr
225 230 235 240

Asn Arg Asp Thr Ile Ser Ile Gly Ala Val Tyr His Tyr Asp Ser Leu
245 250 255

Ile Glu Lys Pro Thr Glu Pro Asn Ala Leu Val Asn Ala Leu Leu Ser
260 265 270

Asn Pro Phe Val Met Glu Leu Ile Lys Asp Glu Val Pro Arg Ile Lys
275 280 285

Glu Asp Tyr Arg Asp Leu Ser Lys Asp Glu Glu Leu Arg Ile Arg Phe
290 295 300

Lys Ser Asn Lys Leu Ile Lys Ser Trp Asn Asp Leu His His Thr Tyr
305 310 315 320

Tyr Ser Pro Ser Ala Val Ala Glu Leu Val Ala Gln Gly Lys Tyr Lys
325 330 335

Ser Arg Glu Glu Ile Lys Asp Lys Ile Asp Ser Leu Tyr Asn Glu Leu
340 345 350

Val Thr Lys Tyr Asn Thr Glu Phe Glu Thr Asn Tyr Val Glu Leu Glu
355 360 365

Tyr Ser Ala Lys Leu Val Pro Asp Gly Lys Arg Cys Arg Met Lys Lys
370 375 380

Pro Tyr Phe Lys Asn Ile Leu Phe Val Gly Asp Ala Ala Gly Arg Gly
385 390 395 400

Ile Phe Leu Gly Pro Arg Ile Glu Gly Leu Asn Val Gly Ile Asp Asp
405 410 415

Ala Val Arg Ala Ala Glu Ala Val Ser Lys Ser Ile Asp Gln Asn Asn
420 425 430

Phe Gln Phe Asp Asn Ile Gly Glu Arg Tyr Thr Lys Ser Val Asp Glu
435 440 445

Ser Pro Tyr Thr Ala Asp Met Ser Arg Ile Asp Ala Asn Tyr Leu Lys
450 455 460

Ala Val Leu Asp Cys Thr Lys Lys Val Pro Lys Asn Thr Leu Gly Phe
465 470 475 480

Lys Tyr Gly Ser Ile Val Lys Leu Met Ser Asn Ser Thr Phe Arg Asn
485 490 495

Val Ser Ile Gly Ile Ala Asn Ser Ile Gly Tyr Lys Arg Leu Leu Pro
500 505 510

Val Ile Glu Ser Asp Lys Thr Tyr Asn Gln Ile Pro Ile Glu Ile Ala
515 520 525

Glu Arg Asn Gly Lys Asp Leu Arg Lys Ser Tyr Ser Ile Glu Ile Pro
530 535 540

Thr Ile Ala Glu Arg Ile Ala Asn Leu Asn Tyr Asn Asp Asp Ser Leu
545 550 555 560

Ser His Ile Lys Val Leu Asn Ser Gln Ser Asp Phe Met Lys Lys Met
565 570 575

Val Gln Leu Cys Pro Thr Lys Cys Tyr Ser Ile Glu Asn Glu Arg Ile
580 585 590

Met Leu Gln His Glu Gly Cys Ile Glu Cys Gly Thr Cys Ala Arg Glu
595 600 605

Thr Glu Trp Arg His Pro Arg Gly Glu Lys Gly Ile Ile Tyr Asn Tyr
610 615 620

Gly
625

<210> 30

<211> 2238

<212> DNA

<213> Crenarchaeote

<400> 30
ttggaagggt cttctctaata acataataat aatctcctatg acattggcac aaataatgat 60
aatgtatgtg aaaataaata cattgacgca aaccccggtt tgcgcacaa acgcggtttt 120
ggtgacgtca acaataacga gaacgataat gatggtggtg ttgacgccgg tgcacccacc 180
acaaagggtg actatggccc tgaaaacgca aacaacgcc ttttgagggt catagacagg 240
gccaatgtga agatagactc ttgcataaac tccgtggccc cgtccgtgat gatagggggt 300
gacgccataa gggagaaaag ggttgacgcg gtcaaaaaca ggggccttaa actgcggtat 360
gtaaccgaaa taacaaagga caacgtcggc tatgtcaagg agatgctctc gttttcggag 420
attaggcacc tggatgggct gaagggaaac tttgaggtgg ccgaccagag ggagtatgtg 480
gctgtcgcca cccttcacgc ggcacagtca ataccocagc tttgttcag caacctccct 540
gagattgcag agcagcagca gtttgtgtt gacagctttt ggggcagggc gttgcccgca 600
gagcacagga taaaggagct ggaggatggg gttgtcatgc ccgtctcact ggtcttctcc 660
aactacaagg acgcggtcca aaggaggtt gaaatgataa gcagggccaa aaggagata 720
ctgataatgt actccacggt taacgcgttc cacctgcagg aaaaaggcgg cacactgcaa 780
ctcttgaagg agatggtgga gcaaacgac agcctgagga tcaacatcct cagccgatg 840
gatgcctcag tgcgagagtc cttgtccttg aggtcctca caaaatacag gcccaacatc 900
caagtccagg acattgcgcc aagcattggc atcaagataa agacactggt tgtggacaga 960
aaggagtgc tggatgatgga gctgatacac gcaaggagg aggtggcaac cgcgcgaatc 1020
ggcttttcaa tctactcaa cagcagacca acggtattgt cttactcgtc catatttgag 1080
gtcctctatg accagagtgt cctgttcag cagcttgacc aaaacgacaa agtcaaaagc 1140
gagttcataa acgtggctgc gcatgagttg cgaacgcaa tcatgccat cctaaacggt 1200
gtggagatac tggaggagaa gcttggcgaa agaaaaacag agtttcagcg ggagcttgac 1260
atgataacaa gaaacgcgtc ccggtgcag aaccttgccg aaagcattct gcaggtgagc 1320
agaatcgaag gcggaagctt tagcctggac atccaaaaaa atgtggatat ccacaacctg 1380
atttcccagg tgatagagga cattgagaaa aaatacgcct acaaggagaa ggcaacaag 1440
gtggcgatag tgtttttgcc atctgacggc aacagaaatg gcgggtactc cagaggtggc 1500
ggcggcgcaa aggcagaagg ggtaaaggca gcggcaggag caaacaggc gcaaaaagag 1560
acgcagcaaa aagagcagtg ggtagaacc gttaatggcc ccaaccacct tttgatgta 1620
gactgtgatc cgcaaaagat aagccaggtt gttttcaacc tgctggacaa cgcaatgaag 1680
ttcaccaatg acggcaagat tgttgtttcc acggcagtga tgggtgagtc ttctcccttc 1740

acttccacct ctcaggaaag tgatacctca aacactgcta cagctggtaa aggcaatggg 1800
ggcagagtgg atagcagcag cgacagcgac aacgggtggtg gtgacaatgg tggtgaccac 1860
atcgggagggc agaaagaagg cgcgggtgcta gtcacagtgc aggacaccgg ggttgggctc 1920
aactccaaaa taagggatca gctgtttcag aaatttgtca caaagtcaaa ccaggggaacc 1980
ggccttggcc tatacctgtc aaggaaaatt gttgaggagc atggtggaaa aatatggttt 2040
gaggagacaa acagcaaggg cggcaacagc agcagcagga acaacactaa agataaagac 2100
gaaggcattg atgaaatact gcaccacctt ggcagtgaag gaaaaatagg cgccacattc 2160
aaatttgtca tacctgtctc cctgccttcc catatgccga caaaagacat gccagaaaaa 2220
aacgatgaag gaaaatga 2238

<210> 31

<211> 745

<212> PRT

<213> Crenarchaeote

<400> 31

Met Glu Gly Ser Ser Leu Ile His Asn Asn Asn Leu His Asp Ile Gly
1 5 10 15

Thr Asn Asn Asp Asn Val Cys Glu Asn Lys Tyr Ile Asp Ala Asn Pro
20 25 30

Gly Leu Ser His Lys Arg Gly Phe Gly Asp Val Asn Asn Asn Glu Asn
35 40 45

Asp Asn Asp Gly Gly Val Asp Ala Gly Ala Pro Thr Thr Lys Val Tyr
50 55 60

Tyr Gly Pro Glu Asn Ala Asn Asn Ala Ile Leu Arg Phe Ile Asp Arg
65 70 75 80

Ala Asn Val Lys Ile Asp Ser Cys Ile Asn Ser Val Ala Pro Ser Val
85 90 95

Met Ile Gly Val Asp Ala Ile Arg Glu Lys Arg Val Asp Ala Val Lys
100 105 110

Asn Arg Gly Leu Lys Leu Arg Tyr Val Thr Glu Ile Thr Lys Asp Asn
115 120 125

Val Gly Tyr Val Lys Glu Met Leu Ser Phe Ser Glu Ile Arg His Leu
130 135 140

Asp Gly Leu Lys Gly Asn Phe Glu Val Ala Asp Gln Arg Glu Tyr Val
145 150 155 160

Ala Val Ala Thr Leu His Ala Ala Gln Ser Ile Pro Gln Leu Leu Phe
165 170 175

Ser Asn Leu Pro Glu Ile Ala Glu Gln Gln Gln Phe Val Phe Asp Ser
180 185 190

Phe Trp Gly Arg Ala Leu Pro Ala Glu His Arg Ile Lys Glu Leu Glu
195 200 205

Asp Gly Val Val Met Pro Val Ser Leu Val Phe Ser Asn Tyr Lys Asp
210 215 220

Ala Val Gln Arg Glu Phe Glu Met Ile Ser Arg Ala Lys Arg Glu Ile
225 230 235 240

Leu Ile Met Tyr Ser Thr Val Asn Ala Phe His Leu Gln Glu Lys Gly
245 250 255

Gly Thr Leu Gln Leu Leu Lys Glu Met Val Glu Gln Asn Asp Ser Leu
260 265 270

Arg Ile Asn Ile Leu Thr Pro Met Asp Ala Ser Val Arg Glu Ser Leu
275 280 285

Ser Leu Arg Leu Leu Thr Lys Tyr Arg Pro Asn Ile Gln Val Gln Asp
290 295 300

Ile Ala Pro Ser Ile Gly Ile Lys Ile Lys Thr Leu Val Val Asp Arg
305 310 315 320

Lys Glu Ser Leu Val Met Glu Leu Ile His Ala Arg Glu Glu Val Ala
325 330 335

Thr Ala Ala Ile Gly Phe Ser Ile Tyr Ser Asn Ser Glu Pro Thr Val
340 345 350

Leu Ser Tyr Ser Ser Ile Phe Glu Val Leu Tyr Asp Gln Ser Val Leu
355 360 365

Phe Gln Gln Leu Asp Gln Asn Asp Lys Val Lys Ser Glu Phe Ile Asn
370 375 380

Val Ala Ala His Glu Leu Arg Thr Pro Ile Met Pro Ile Leu Asn Gly
385 390 395 400

Val Glu Ile Leu Glu Glu Lys Leu Gly Glu Arg Lys Thr Glu Phe Gln
405 410 415

Arg Glu Leu Asp Met Ile Thr Arg Asn Ala Ser Arg Leu Gln Asn Leu
420 425 430

Ala Glu Ser Ile Leu Gln Val Ser Arg Ile Glu Ser Gly Ser Phe Ser
435 440 445

Leu Asp Ile Gln Lys Asn Val Asp Ile His Asn Leu Ile Ser Gln Val
450 455 460

Ile Glu Asp Ile Glu Lys Lys Tyr Ala Tyr Lys Glu Lys Ala Asn Lys
465 470 475 480

Val Ala Ile Val Phe Leu Pro Ser Asp Gly Asn Arg Asn Gly Gly Tyr
485 490 495

Ser Arg Gly Gly Gly Gly Ala Lys Ala Glu Gly Val Lys Ala Ala Ala
500 505 510

Gly Ala Lys Gln Ala Gln Lys Glu Thr Gln Gln Lys Glu Gln Trp Val
515 520 525

Glu Pro Val Asn Gly Pro Asn His Leu Leu Tyr Val Asp Cys Asp Pro
530 535 540

Gln Lys Ile Ser Gln Val Val Phe Asn Leu Leu Asp Asn Ala Met Lys
545 550 555 560

Phe Thr Asn Asp Gly Lys Ile Val Val Ser Thr Ala Val Met Gly Glu
565 570 575

Ser Ser Pro Phe Thr Ser Thr Ser Gln Glu Ser Asp Thr Ser Asn Thr
580 585 590

Ala Thr Ala Gly Lys Gly Asn Gly Gly Arg Val Asp Ser Ser Ser Asp
595 600 605

Ser Asp Asn Gly Gly Gly Asp Asn Gly Gly Asp His Ile Gly Arg Gln
610 615 620

Lys Glu Gly Ala Val Leu Val Thr Val Gln Asp Thr Gly Val Gly Leu
625 630 635 640

Asn Ser Lys Ile Arg Asp Gln Leu Phe Gln Lys Phe Val Thr Lys Ser
645 650 655

Asn Gln Gly Thr Gly Leu Gly Leu Tyr Leu Ser Arg Lys Ile Val Glu
660 665 670

Glu His Gly Gly Lys Ile Trp Phe Glu Glu Thr Asn Ser Lys Gly Gly
675 680 685

Asn Ser Ser Ser Arg Asn Asn Thr Lys Asp Lys Asp Glu Gly Ile Asp
690 695 700

Glu Ile Leu His His Leu Gly Ser Glu Gly Lys Ile Gly Ala Thr Phe
705 710 715 720

Lys Phe Val Ile Pro Val Ser Leu Pro Ser His Met Pro Thr Lys Asp
725 730 735

Met Pro Glu Lys Asn Asp Glu Gly Lys
740 745

<210> 32

<211> 519

<212> DNA

<213> Crenarchaeote

<400> 32
ttgcaaagca gtcacatcttc taaaataatc acaatttgca gaatgccgtc acttcatctt 60
gttgcatatg gtttaatttt ggatattttc gaaagcccaa tcacaagggt aaacggtaga 120
acaagtcact tgattattaa aatatatcca catatggata acaatacaag gatgagttct 180
ttagcaatog agtttttttt atcccttttt tcaataacgt tactttctaa aagaatatac 240
caaccagtga aatcaaagtc atatacctac catgacaagc atccatttca gtacaagatg 300

gaggattatg caaaccacaa caaaattgta gactataaaa actgcttact tttttttcaa 360
gtatcgatgt tacaaaaaaa taaaataatt aggattcggg ttccagggtt gttttataca 420
ggtggctgga tttccctcac actaaagttt ttgatatcca catcatttgc accatcccac 480
ctgaaagtag caatggggcc tccccaggat ataatctga 519

<210> 33

<211> 172

<212> PRT

<213> Crenarchaeote

<400> 33

Met Gln Ser Ser His Leu Ser Lys Ile Ile Thr Ile Cys Arg Met Pro
1 5 10 15

Ser Leu His Leu Val Ala Tyr Gly Leu Ile Leu Asp Ile Phe Glu Ser
20 25 30

Pro Ile Thr Arg Leu Asn Gly Arg Thr Ser His Leu Ile Ile Lys Ile
35 40 45

Tyr Pro His Met Asp Asn Asn Thr Arg Met Ser Ser Leu Ala Ile Glu
50 55 60

Phe Phe Leu Ser Leu Phe Ser Ile Thr Leu Leu Ser Lys Arg Ile Tyr
65 70 75 80

Gln Pro Val Lys Ser Lys Ser Tyr Thr Tyr His Asp Lys His Pro Phe
85 90 95

Gln Tyr Lys Met Glu Asp Tyr Ala Asn His Asn Lys Ile Val Asp Tyr
100 105 110

Lys Asn Cys Leu Leu Phe Phe Gln Val Ser Met Leu Gln Lys Asn Lys
115 120 125

Ile Ile Arg Ile Arg Val Pro Gly Leu Phe Tyr Thr Gly Gly Trp Ile
130 135 140

Ser Leu Thr Leu Lys Phe Leu Ile Ser Thr Ser Phe Ala Pro Ser His
145 150 155 160

Leu Lys Val Ala Met Gly Pro Pro Gln Asp Ile Ile.
165 170

<210> 34

<211> 1008

<212> DNA

<213> Crenarchaeote

<400> 34

```
acactgctaa caagcggagg cgcacctgtg cccgcaaccg taagcatgaa ttcaccaacc      60
aacacagttg cgaccctaaa cccctctgca gatttaacac ctggtgccac atacactgcc      120
agaataacaa ccggtgctac ggatgcgact ggtgtcccat tggctgccga caaggtatgg      180
acgttctctg ttgcccttc tggaggcgga gggacattag accagtttgg gataaccag      240
atttaccocg ataaggcggg tgggtggagaa aaatggttta tgaatatgca aaaccogaac      300
aatgatccac gaacaaaccc acctgacatg gacctaaacc cagatggcag ttggaacgtt      360
aatgacgatc aggtcagata taacgtgttt acatcatcag ggtaccatcc agaggatatt      420
gagacttacg atcactcggg actcgcaaca caaggataca tgcagtatcc aaatgattgg      480
aagaatgtgg aaatgacggg tattgtaagg gttaatagtg gagatgattc tgaaaatttc      540
gcttggtatg acaggggtgg taggcactat gatggcgaag gatgagagg ctcagcatat      600
aaagcagatc tattctatga tggaagggtt aggcttgcaa aagagcagtg gcatgtctcc      660
tatgtgtttt caagcactac cacaccctca ccttcggcgt ctagttttga tagattcatt      720
ggatttaaag ccatgatcta taaccaccaa ttggtggtg gtgagaccgt agtcaccact      780
gaaatatggg tagacagaaa cccggatagc ccgactctga agaacaattg gcaaaaggta      840
tacacattca ctgactcagg tgggtttgga aatgatggtg aagagtgtgg tggtagccg      900
gatcagatta tctctgggg aggccccatt gctactttca ggtgggatgg tgcaaatgat      960
gtggatatca aaaactttag tgtgagggaa atccagccac ctgtataa      1008
```

<210> 35

<211> 335

<212> PRT

<213> Crenarchaeote

<400> 35

Thr Leu Leu Thr Ser Gly Gly Ala Pro Val Pro Ala Thr Val Ser Met
1 5 10 15

Asn Ser Pro Thr Asn Thr Val Ala Thr Leu Asn Pro Ser Ala Asp Leu
20 25 30

Thr Pro Gly Ala Thr Tyr Thr Ala Arg Ile Thr Thr Gly Ala Thr Asp
35 40 45

Ala Thr Gly Val Pro Leu Ala Ala Asp Lys Val Trp Thr Phe Ser Val
50 55 60

Ala Pro Ser Gly Gly Gly Gly Thr Leu Asp Gln Phe Gly Ile Thr Gln
65 70 75 80

Ile Tyr Pro Asp Lys Ala Gly Gly Gly Glu Lys Trp Phe Met Asn Met
85 90 95

Gln Asn Pro Asn Asn Asp Pro Arg Thr Asn Pro Pro Asp Met Asp Leu
100 105 110

Asn Pro Asp Gly Ser Trp Asn Val Asn Asp Asp Gln Val Arg Tyr Asn
115 120 125

Val Phe Thr Ser Ser Gly Tyr His Pro Glu Asp Ile Glu Thr Tyr Asp
130 135 140

His Ser Val Leu Ala Thr Gln Gly Tyr Met Gln Tyr Pro Asn Asp Trp
145 150 155 160

Lys Asn Val Glu Met Thr Gly Ile Val Arg Val Asn Ser Gly Asp Asp
165 170 175

Ser Glu Asn Phe Ala Trp Tyr Asp Arg Gly Gly Arg His Tyr Asp Gly
180 185 190

Glu Gly Cys Glu Gly Ser Ala Tyr Lys Ala Asp Leu Phe Tyr Asp Gly
195 200 205

Arg Val Arg Leu Ala Lys Glu Gln Trp His Val Ser Tyr Val Phe Ser
210 215 220

Ser Thr Thr Thr Pro Ser Pro Ser Ala Ser Ser Phe Asp Arg Phe Ile
225 230 235 240

Gly Phe Lys Ala Met Ile Tyr Asn His Gln Leu Ala Gly Gly Glu Thr
245 250 255

Val Val Thr Thr Glu Ile Trp Val Asp Arg Asn Pro Asp Ser Pro Thr
260 265 270

Leu Lys Asn Asn Trp Gln Lys Val Tyr Thr Phe Thr Asp Ser Gly Gly
275 280 285

Phe Gly Asn Asp Gly Glu Glu Cys Gly Gly Glu Pro Asp Gln Ile Ile
290 295 300

Ser Trp Gly Gly Pro Ile Ala Thr Phe Arg Trp Asp Gly Ala Asn Asp
305 310 315 320

Val Asp Ile Lys Asn Phe Ser Val Arg Glu Ile Gln Pro Pro Val
325 330 335

1 / 13




Figure 1A




Figure 1B

2 / 13




Figure 2




Figure 3

3 / 13




Figure 4

Figure 5 B





Figure 5 A



5 / 13




Figure 6

6 / 13

Figure 7




Figure 8 A





Figure 8B



7 / 13

Figure 9 A





Figure 9 B



8 / 13



Figure 10




Figure 11

9 / 13

Figure 12





Figure 13



10 / 13




Figure 14 A

11 / 13




Figure 14 B

12 / 13




Figure 14 C

13 / 13




Figure 14 D

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.